

PERFECTION IN MOTIVIC HOMOTOPY THEORY

ELDEN ELMANTO AND ADEEL A. KHAN

ABSTRACT. We prove a topological invariance statement for the Morel–Voevodsky motivic homotopy category, up to inverting the exponential characteristic p of the base field. This implies in particular that $\mathbf{SH}[\frac{1}{p}]$ is invariant under passing to perfections. Among other applications we prove Grothendieck–Verdier duality in this context.

CONTENTS

1. Introduction	1
2. Topological invariance	2
2.1. Main result and corollaries	2
2.2. Proof of Theorem 2.1.1	4
3. Applications	6
3.1. Duality	6
3.2. Removal of perfectness hypotheses	6
References	8

1. INTRODUCTION

Let S be a scheme and denote by $S_{\text{ét}}$ its small étale topos. The starting point for this note is Grothendieck’s “équivalence remarquable de catégories” [Gro67, Théorème 18.1.2], which asserts that for any nil-immersion $f : S_0 \hookrightarrow S$, there is an induced equivalence

$$f^* : S_{\text{ét}} \rightarrow (S_0)_{\text{ét}}.$$

In fact, Grothendieck further generalized this to a *topological invariance* statement for the small étale topos: for any universal homeomorphism of schemes $f : T \rightarrow S$, the functor $f^* : S_{\text{ét}} \rightarrow T_{\text{ét}}$ is an equivalence (see [GR71, Exposé IX, Théorème 4.10], [AGV72, Exposé VIII, Théorème 1.1]).

The large étale topos fails to satisfy nil-invariance. An observation of Morel and Voevodsky [MV99] was that this failure can be repaired by working in the setting of \mathbf{A}^1 -invariant sheaves. Indeed, it is a consequence of the Morel–Voevodsky localization theorem that the stable motivic homotopy category \mathbf{SH} satisfies nil-invariance (see e.g. [CD12, Proposition 2.3.6(1)]). However, the topological invariance property still fails, at least in positive characteristic (see Remark 2.1.9). Our goal in this paper is to show that topological invariance is in fact true for \mathbf{SH} , after inverting the exponential characteristic of the base field (Theorem 2.1.1). This also recovers the analogous result for mixed motives as proven in [CD15].

A particularly useful consequence of topological invariance is that, for any scheme S of characteristic p , $\mathbf{SH}(S)[\frac{1}{p}]$ is invariant under passing to the perfection S_{perf} (Corollary 2.1.4). This allows us to remove perfectness hypotheses on the base field in many results, see §3.2. It also yields a Grothendieck–Verdier duality statement, following ideas of Cisinski–Déglise [CD15] (Theorem 3.1.1).

Date: December 19, 2018.

1.1. Conventions.

1.1.1. All schemes will implicitly be assumed to be quasi-compact quasi-separated.

Recall that a morphism of schemes $f : X \rightarrow Y$ is a *universal homeomorphism* if it induces a homeomorphism on underlying topological spaces after any base change, or equivalently, if it is integral, universally injective, and surjective [Gro67, Corollary 18.12.11].

1.1.2. If S is a scheme of characteristic $p > 0$, i.e., an \mathbf{F}_p -scheme, we write $F_S : S \rightarrow S$ for the Frobenius endomorphism [Gro77, Exposé XIV=XV]. Recall that S is *perfect* if the Frobenius $F_S : S \rightarrow S$ is an isomorphism. Any \mathbf{F}_p -scheme S admits a *perfection* S_{perf} , defined as the limit of the tower

$$\dots \xrightarrow{F_S} S \xrightarrow{F_S} S,$$

see [BS17, Section 3].

1.1.3. Given a scheme S , we denote by $\mathbf{SH}(S)$ the stable ∞ -category of motivic spectra over S . We will use the language of six operations, see [Hoy14, Appendix C] or [Kha16] for the non-noetherian setting. Any motivic spectrum $E \in \mathbf{SH}(S)$ represents a cohomology theory on S -schemes, given by the formula

$$E(X/S, \xi) = \text{Maps}_{\mathbf{SH}(S)}(\mathbf{1}_S, f_* \Sigma^\xi f^*(E))$$

for any morphism $f : X \rightarrow S$ and any K -theory class $\xi \in K(X)$. Similarly, there is a Borel–Moore homology theory

$$E^{\text{BM}}(X/S, \xi) = \text{Maps}_{\mathbf{SH}(S)}(\mathbf{1}_S, f_* \Sigma^{-\xi} f^!(E)).$$

We refer to [DJK18] for details.

1.2. Acknowledgements. We would like to thank Denis-Charles Cisinski, Frédéric Déglise, Marc Hoyois, and Fangzhou Jin for useful suggestions and conversations on the subject of this paper. We would also like to thank the Center for Symmetry and Deformation at the University of Copenhagen for supporting AK’s visit during which this paper was conceived. EE would like to thank a couple of cheeky colleagues for suggesting the title.

2. TOPOLOGICAL INVARIANCE

2.1. Main result and corollaries.

Theorem 2.1.1. *Let S be a scheme of exponential characteristic p . Then for any universal homeomorphism $f : T \rightarrow S$, the functor*

$$f^* : \mathbf{SH}(S)[\frac{1}{p}] \rightarrow \mathbf{SH}(T)[\frac{1}{p}]$$

is an equivalence.

Corollary 2.1.2. *Let S be a scheme of exponential characteristic p . Then for any universal homeomorphism $f : T \rightarrow S$, there is an equivalence of functors*

$$f^* \simeq f^! : \mathbf{SH}(S)[\frac{1}{p}] \rightarrow \mathbf{SH}(T)[\frac{1}{p}].$$

Proof. Since f^* is an equivalence, its quasi-inverse is given by $f_* \simeq f_!$. Hence we have an equivalence of the left and right adjoints of this latter functor. \square

Corollary 2.1.3. *For every scheme S of characteristic $p > 0$, the absolute Frobenius induces an equivalence*

$$F_S^* : \mathbf{SH}(S)[\frac{1}{p}] \rightarrow \mathbf{SH}(S)[\frac{1}{p}].$$

Corollary 2.1.4. *For every scheme S of characteristic $p > 0$, the canonical morphism $S_{\text{perf}} \rightarrow S$ induces an equivalence*

$$\mathbf{SH}(S)[\frac{1}{p}] \rightarrow \mathbf{SH}(S_{\text{perf}})[\frac{1}{p}].$$

Proof. Follows from Corollary 2.1.3 in view of continuity of \mathbf{SH} [Hoy14, Proposition C.12(4)]. \square

2.1.5. At the level of cohomology and Borel–Moore homology, we have the following reformulation:

Corollary 2.1.6. *Let S be a scheme of exponential characteristic p . Let $E \in \mathbf{SH}(S)$ be a motivic spectrum over S . Then we have:*

(i) *For any universal homeomorphism $f : X \rightarrow Y$ of S -schemes, the induced maps*

$$\begin{aligned} f^* : E(Y, \xi)[\frac{1}{p}] &\rightarrow E(X, f^*(\xi))[\frac{1}{p}] \\ f_* : E^{\text{BM}}(X, f^*(\xi))[\frac{1}{p}] &\rightarrow E^{\text{BM}}(Y, \xi)[\frac{1}{p}] \end{aligned}$$

are equivalences for every $\xi \in K(Y)$.

(ii) *The canonical morphism $f : S_{\text{perf}} \rightarrow S$ induces equivalences*

$$\begin{aligned} f^* : E(S, \xi)[\frac{1}{p}] &\rightarrow E(S_{\text{perf}}, f^*(\xi))[\frac{1}{p}] \\ f_* : E^{\text{BM}}(S_{\text{perf}}, f^*(\xi))[\frac{1}{p}] &\rightarrow E^{\text{BM}}(S, \xi)[\frac{1}{p}] \end{aligned}$$

for every $\xi \in K(S)$.

Proof. Note that we have canonical identifications

$$E(T, \psi)[\frac{1}{p}] \simeq \text{Maps}_{\mathbf{SH}(S)}(\mathbf{1}_S, (\pi_T)_* \Sigma^\psi \pi_T^*(E))[\frac{1}{p}] \simeq \text{Maps}_{\mathbf{SH}(S)[\frac{1}{p}]}(\mathbf{1}_S, (\pi_T)_* \Sigma^\psi \pi_T^*(E))$$

for every $\pi_T : T \rightarrow S$ and $\psi \in K(T)$. Therefore the map on cohomology spaces is induced from the natural transformation

$$(\pi_Y)_* \Sigma^\xi \pi_Y^* \rightarrow (\pi_Y)_* \Sigma^\xi f_* f^* \pi_Y^* \simeq (\pi_Y)_* f_* \Sigma^{f^*(\xi)} f^* \pi_Y^* \simeq (\pi_X)_* \Sigma^{f^*(\xi)} \pi_X^*.$$

For f as in (i) (resp. (ii)), the unit map $\text{id} \rightarrow f_* f^*$ is invertible after inverting p by Theorem 2.1.1 (resp. by Corollary 2.1.4), whence the claim. The proof for Borel–Moore homology is similar, using the fact that the co-unit map $f_* f^! \rightarrow \text{id}$ is also invertible in both cases (after inverting p). \square

Remark 2.1.7. Corollary 2.1.6 also holds for the compactly supported variants (cohomology with compact support and relative homology), with the same proofs.

Example 2.1.8. Let $\text{KGL} \in \mathbf{SH}(\mathbf{F}_p)$ denote the homotopy invariant K-theory spectrum over $\text{Spec}(\mathbf{F}_p)$. For every (possibly singular and non-noetherian) \mathbf{F}_p -scheme S , we have functorial equivalences

$$\text{KGL}(S, 0)[\frac{1}{p}] \simeq K(S)[\frac{1}{p}]$$

by [Cis13, Theorem 2.20] and [TT90, Exercise 9.11(h)]. Under these identifications, Corollary 2.1.6 recovers in particular the recent observation of Kelly and Morrow [KM18, Lemma 4.1] that the canonical map

$$K(S)[\frac{1}{p}] \rightarrow K(S_{\text{perf}})[\frac{1}{p}]$$

is an equivalence.

Remark 2.1.9. If $p > 1$, then Theorem 2.1.1 is *false* before inverting p . Indeed, if k is a non-perfect field, then the functor $F^* : \mathbf{SH}(k) \rightarrow \mathbf{SH}(k)$ is not an equivalence. Supposing the contrary, it would follow that the induced map

$$F^* : K(k) \rightarrow K(k)$$

is an equivalence (as in Corollary 2.1.6). However, as the Frobenius map $k \rightarrow k$ is by assumption *not* an isomorphism, the map

$$k^\times = K_1(k) \rightarrow k^\times = K_1(k),$$

which is additively given by multiplication by p , is not an isomorphism.

2.2. Proof of Theorem 2.1.1.

Notation 2.2.1. Given a unit $a \in \Gamma(S, \mathcal{O}_S)^\times$, we write $\langle a \rangle$ for the induced point of $\Omega K(S)$. For an integer $n \geq 0$, we write n_ϵ for the formal sum

$$n_\epsilon = 1 + \langle -1 \rangle + 1 + \cdots$$

which consists of n terms. We may also regard n_ϵ as an automorphism of the identity functor $\mathrm{id}_{\mathbf{SH}(S)}$, via the canonical map $\Omega K(S) \rightarrow \mathrm{Aut}(\mathrm{id}_{\mathbf{SH}(S)})$.

We are grateful to Marc Hoyois for suggesting the following re-interpretation of [EHK⁺18b, Proposition B.1.4].

Proposition 2.2.2. *Let S be a scheme, $P \in \Gamma(S, \mathcal{O}_S)[x]$ a monic polynomial of degree d , and $T \subset \mathbf{A}_S^1$ the closed subscheme cut out by P . If $f : T \rightarrow S$ denotes the canonical morphism, then there exists a canonical natural transformation*

$$\mathrm{tr}_f : f_* f^* \rightarrow \mathrm{id}_{\mathbf{SH}(S)}$$

such that the composites

$$\mathrm{id} \xrightarrow{\mathrm{unit}} f_* f^* \xrightarrow{\mathrm{tr}_f} \mathrm{id}, \quad f_* f^* \xrightarrow{\mathrm{tr}_f} \mathrm{id} \xrightarrow{\mathrm{unit}} f_* f^*$$

are homotopic to d_ϵ and $f_* * d_\epsilon * f^*$, respectively.

Proof. Note that $f : T \rightarrow S$ is finite and syntomic. The conormal sheaf $\mathcal{N}_{T/\mathbf{A}_S^1} \simeq (P)/(P^2)$ is free of rank 1, and the generator P induces a canonical trivialization $\tau : \mathbf{L}_f \simeq 0$ in $K(T)$. Therefore, the trace transformation tr_f of [DJK18] induces a canonical natural transformation

$$f_* f^* \simeq f_* \Sigma^{\mathbf{L}_f} f^* \xrightarrow{\mathrm{tr}_f} \mathrm{id}$$

which we denote again by tr_f . The claim can be reformulated in cohomological terms as the assertion that, for every $E \in \mathbf{SH}(S)$, the composites

$$(2.2.3) \quad E(S, 0) \xrightarrow{f^*} E(T, 0) \simeq E(T, \langle \mathbf{L}_f \rangle) \xrightarrow{f_!} E(S, 0),$$

$$(2.2.4) \quad E(T, 0) \simeq E(T, \langle \mathbf{L}_f \rangle) \xrightarrow{f_!} E(S, 0) \xrightarrow{f^*} E(T, 0)$$

are homotopic to multiplication by d_ϵ .

By [EHK⁺18a, Theorem 3.2.11] the first composite is homotopic to the transfer map induced by the framed correspondence

$$\begin{array}{ccc} & T & \\ f, \tau \swarrow & & \searrow f \\ S & & S. \end{array}$$

Therefore the claim follows from [EHK⁺18b, Proposition B.1.4] applied to $E(-, 0)$, viewed as a presheaf on $\text{Corr}^{\text{fr}}(\text{Sm}/S)$. Similarly, the second composite is identified, by the transverse base change property of the trace transformation tr_f [DJK18, Proposition 2.3.6], with

$$E(\mathbf{T}, 0) \xrightarrow{\pi_2^*} E(\mathbf{T} \times_S \mathbf{T}, 0) \simeq E(\mathbf{T} \times_S \mathbf{T}, \langle L_f \rangle) \xrightarrow{(\pi_1)_!} E(\mathbf{T}, 0),$$

where π_1 and π_2 are the first and second projections of $\mathbf{T} \times_S \mathbf{T}$, respectively. As above, this is identified with the transfer map induced by the framed correspondence

$$\begin{array}{ccc} & \mathbf{T} \times_S \mathbf{T} & \\ \pi_1, \pi_2^*(\tau) \swarrow & & \searrow \pi_2 \\ \mathbf{T} & & \mathbf{T} \end{array}$$

so the claim follows by another application of [EHK⁺18b, Proposition B.1.4]. \square

Lemma 2.2.5. *Let S be the spectrum of a field k of exponential characteristic p . Then for any power q of p , the canonical map*

$$\text{End}_{\mathbf{SH}(S)}(\mathbf{1}_S) \rightarrow \text{End}_{\mathbf{SH}(S)}(\mathbf{1}_S)[\frac{1}{p}] \simeq \text{End}_{\mathbf{SH}(S)[\frac{1}{p}]}(\mathbf{1}_S)$$

sends $q_\epsilon \in \text{End}_{\mathbf{SH}(S)}(\mathbf{1}_S)$ to a unit.

Proof. We only need to consider the case $p > 1$. Using Morel's identification $\text{End}(\mathbf{1}_S) \simeq \text{GW}(k)$ [Mor04], which has been extended in [BH18, Lemma 10.12], it will suffice to show that the induced element $q_\epsilon \in \text{GW}(k)[\frac{1}{p}]$ is invertible. In view of the cartesian square

$$\begin{array}{ccc} \text{GW}(k) & \xrightarrow{\dim} & \mathbf{Z} \\ \downarrow & & \downarrow \\ \text{W}(k) & \longrightarrow & \mathbf{Z}/2 \end{array}$$

as in [Mor12, (3.1)] (cf. [Bac18, Lemma 17], [KK82, Lemma 1.16]), it will in fact suffice to only check invertibility in $\mathbf{Z}[\frac{1}{p}]$ and in $\text{W}(k)[\frac{1}{p}]$. The former is obvious. For the latter, we first assume that p is odd. In this case, we note that $d_\epsilon = \frac{d-1}{2}h + 1$ and thus q_ϵ is invertible in $\text{W}(k)$ (without inverting p). When $p = 2$, -1 is trivially a sum of squares in k , so the Witt ring is 2-torsion [MH73, Theorem III.3.6] and the claim follows. \square

Lemma 2.2.6. *Let S be a scheme of exponential characteristic p . If $f : \mathbf{T} \rightarrow S$ is a universal homeomorphism, then the functor*

$$f^* : \mathbf{SH}(S)[\frac{1}{p}] \rightarrow \mathbf{SH}(\mathbf{T})[\frac{1}{p}]$$

is fully faithful.

Proof. The claim is that the unit map $\text{id} \rightarrow f_* f^*$ is invertible after inverting p . By continuity [Hoy14, Proposition C.12(4)] and proper base change, we may use a noetherian approximation argument [TT90, Theorem C.9] to assume that S is noetherian and of finite dimension. Then using [BH18, Proposition A.3] (and proper base change again), we may assume that S is a henselian local scheme; we denote its closed point by $i : \{s\} \rightarrow S$ and the complement by $j : U \rightarrow S$. By the localization theorem, the pair of functors (i^*, j^*) is jointly conservative (see [CD12, Section 2.3]). Since U has dimension strictly lower than that of S , we can argue by induction on the dimension on S to reduce to the case where $S = \{s\}$, i.e., where S is the spectrum of a field k . Since f is radicial, it is then induced by a purely inseparable field extension $k \subset K$. In characteristic zero ($p = 1$), we are already done. Otherwise, by using continuity again, we may assume that the extension $k \subset K$ is finite, i.e., that $K = k(\alpha)$ with $\alpha^q \in k$ for q some power of the prime p . Now the claim follows from Proposition 2.2.2 and Lemma 2.2.5. \square

Theorem 2.1.1 now follows from Lemma 2.2.6 by an argument of Cisinski–Déglise [CD12, Proposition 2.1.9], which we reproduce here for the reader’s convenience.

Proof of Theorem 2.1.1. By Lemma 2.2.6 it will suffice to show that the co-unit $f^*f_* \rightarrow \text{id}$ is invertible. By the proper base change formula ([Ayo08, Corollary 1.7.18], [CD12, Proposition 2.3.11]), this is identified with the natural transformation $(\pi_2)_*(\pi_1)^* \rightarrow \text{id}$, where π_1 and π_2 are the respective projections $T \times_S T \rightarrow T$. Since f is a universal homeomorphism, its diagonal $\Delta : T \rightarrow T \times_S T$ is a nilpotent closed immersion. Then by the localization theorem (cf. [CD12, Proposition 2.3.6(1)]), Δ^* is an equivalence. Since π_1 and π_2 are retractions of Δ it follows that we have canonical identifications $\Delta^* \simeq (\pi_\varepsilon)_*$ and $\Delta_* \simeq (\pi_\varepsilon)^*$ for each $\varepsilon \in \{1, 2\}$. In particular, the natural transformation $(\pi_2)_*(\pi_1)^* \rightarrow \text{id}$ is identified with the co-unit $\Delta^*\Delta_* \rightarrow \text{id}$, which is invertible. \square

3. APPLICATIONS

3.1. Duality. Let S be a scheme that is locally of finite type over a field k of exponential characteristic p . The structural morphism $\pi : S \rightarrow \text{Spec}(k)$ determines a *duality functor* defined by

$$D_S(E) = \underline{\text{Hom}}(E, \pi^!(\mathbf{1}_k)).$$

To justify this name, we must show that the object $\pi^!(\mathbf{1}_k)$ is *dualizing*. That is:

Theorem 3.1.1. *For any compact object $E \in \mathbf{SH}(S)$, the canonical map*

$$E \rightarrow D_S(D_S(E))$$

is an equivalence in $\mathbf{SH}(S)[\frac{1}{p}]$.

Remark 3.1.2. As remarked in [CD15, Remark 7.4], Theorem 3.1.1 implies the formalism of Grothendieck–Verdier duality for $\mathbf{SH}[\frac{1}{p}]$, for locally of finite type k -schemes. In particular, this gives an improvement of [BD15, Theorem 2.4.8].

Theorem 3.1.1 follows from the following statement, analogous to [CD15, Proposition 7.2].

Proposition 3.1.3. *The full subcategory of compact objects in $\mathbf{SH}(S)[\frac{1}{p}]$ is generated as a thick subcategory by objects of the form $f_!(\mathbf{1})(n)$, where $f : X \rightarrow S$ proper, X is smooth over a purely inseparable extension of k , and $n \in \mathbf{Z}$ is an integer.*

Proof. If k is perfect, the statement is [BD15, Corollary 2.4.7]. In general, the morphism $\varphi : S_{\text{perf}} \simeq S \times_{\text{Spec}(k)} \text{Spec}(k_{\text{perf}}) \rightarrow S$ induces an equivalence $\varphi^* : \mathbf{SH}(S) \rightarrow \mathbf{SH}(S_{\text{perf}})$ by Corollary 2.1.4. If $f : X \rightarrow S_{\text{perf}}$ is a proper morphism with X smooth over k_{perf} , then the composite $X \rightarrow S_{\text{perf}} \rightarrow S$ is as in the statement, so we conclude. \square

Proof of Theorem 3.1.1. As in the proof of [CD15, Theorem 7.3], this follows immediately follows from Proposition 3.1.3, and Ayoub’s purity theorem for smooth morphisms [Ayo08, Section 1.6]. \square

3.2. Removal of perfectness hypotheses. Corollary 2.1.4 allows us to immediately drop perfectness hypotheses in many known results, at least after inverting the exponential characteristic. Some examples are listed below.

Theorem 3.2.1. *Let S be the spectrum of a field k of exponential characteristic p . For any smooth S -scheme X , the suspension spectrum $\Sigma_{\mp}^{\infty}(X)$ is strongly dualizable in $\mathbf{SH}(S)[\frac{1}{p}]$. In particular, $\mathbf{SH}(S)[\frac{1}{p}]$ is generated under colimits by the strongly dualizable objects.*

Indeed, we can use Corollary 2.1.4 to reduce the case where k is perfect, which is due to Riou, see [LYZ13, Corollary B.2].

Remark 3.2.2. Proposition 2.2.2 gives the following refinement of [LYZ13, Lemma B.3]. Suppose $f : Y \rightarrow X$ is a finite étale morphism of degree d between smooth connected k -schemes. Then, up to replacing X by a dense open subset $U \subseteq X$ and f by its base change $f_U : Y_U \rightarrow U$, there are isomorphisms of natural transformations

$$\begin{aligned} (\mathrm{id} \xrightarrow{\mathrm{unit}} f_* f^* \simeq f_{\#} f^* \xrightarrow{\mathrm{counit}} \mathrm{id}) &\simeq d_{\epsilon}, \\ (f_* f^* \simeq f_{\#} f^* \xrightarrow{\mathrm{counit}} \mathrm{id} \xrightarrow{\mathrm{unit}} f_* f^* \mathrm{id}) &\simeq f_* * d_{\epsilon} * f^*, \end{aligned}$$

where d_{ϵ} is as in Notation 2.2.1. To prove this, note there are canonical identifications $\Sigma^{L_f} \simeq \mathrm{id}$ and $f_* \simeq f_! \simeq f_{\#}$ since f is finite and étale, and the composite

$$f_* f^* \simeq f_{\#} f^* \xrightarrow{\mathrm{counit}} \mathrm{id}$$

is canonically homotopic to the trace transformation $\mathrm{tr}_f : f_* f^* \simeq f_* \Sigma^{L_f} f^* \rightarrow \mathrm{id}$. Replacing X by its generic point, we may assume that $X = \mathrm{Spec}(k)$. Then $Y = \mathrm{Spec}(K)$ with K/k a finite separable field extension, so by the primitive element theorem we are now in the situation of Proposition 2.2.2.

3.2.3. We also have the following variant of Bachmann's conservativity theorem [Bac18].

Theorem 3.2.4. *Let k be a field with finite 2-étale cohomological dimension and exponential characteristic p . Then the functor*

$$\mathbf{SH}(k)[\frac{1}{p}] \rightarrow \mathbf{DM}(k; \mathbf{Z}[\frac{1}{p}]),$$

is conservative on compact objects

Proof. Using Corollary 2.1.4 and the analogous result for mixed motives [CD15, Lemma 3.15], we may replace k by its perfection. Then the result is proven in [Bac18, Theorem 16]. \square

Remark 3.2.5. Using Theorem 3.2.4 we can deduce the Pic-injectivity result of [Bac18, Theorem 18]. This extends Bachmann's results on Po Hu's conjecture on invertibility of the the suspension spectra of affine quadrics to imperfect fields; see *loc. cit* for details.

3.2.6. We can similarly extend the recognition principle for infinite loop spaces [EHK⁺18b, Theorem 3.5.13] to non-perfect fields.

Theorem 3.2.7. *Let k be a field of exponential characteristic p . Then there are canonical equivalences of symmetric monoidal ∞ -categories*

$$\begin{aligned} \mathbf{H}^{\mathrm{fr}}(k)^{\mathrm{gp}}[\frac{1}{p}] &\simeq \mathbf{SH}^{\mathrm{veff}}(k)[\frac{1}{p}], \\ \mathbf{SH}^{\mathrm{S}^1, \mathrm{fr}}(k)[\frac{1}{p}] &\simeq \mathbf{SH}^{\mathrm{eff}}(k)[\frac{1}{p}]. \end{aligned}$$

Remark 3.2.8. Theorem 3.2.7 also implies cancellation in the sense of [EHK⁺18b, Theorem 3.5.8] for non-perfect fields, after inverting the exponential characteristic.

Lemma 3.2.9. *Let k be a field of exponential characteristic p . Then the morphism $f : \mathrm{Spec}(k_{\mathrm{perf}}) \rightarrow \mathrm{Spec}(k)$ induces an equivalence*

$$f^* : \mathbf{H}^{\mathrm{fr}}(k)^{\mathrm{gp}}[\frac{1}{p}] \rightarrow \mathbf{H}^{\mathrm{fr}}(k_{\mathrm{perf}})^{\mathrm{gp}}[\frac{1}{p}]$$

of symmetric monoidal ∞ -categories.

Proof. We first show fully faithfulness. By continuity, it suffices to show that the Frobenius induces fully faithful functors $F^* : \mathbf{H}^{\mathrm{fr}}(k)^{\mathrm{gp}}[\frac{1}{p}] \rightarrow \mathbf{H}^{\mathrm{fr}}(k)^{\mathrm{gp}}[\frac{1}{p}]$, i.e., that the counit map $\mathrm{id} \rightarrow$

F_*F^* is an equivalence after inverting p . For this it suffices to show that, for every grouplike framed motivic space \mathcal{F} and every smooth k -scheme X , the induced map of spaces

$$F^* : \mathcal{F}(X)_{[\frac{1}{p}]} \rightarrow \mathcal{F}(F^{-1}(X))_{[\frac{1}{p}]}$$

is invertible. As in the proof of Proposition 2.2.2, the two composites (2.2.3) and (2.2.4) are now part of the structure of \mathcal{F} , so that we can appeal directly to [EHK⁺18b, Proposition B.1.4]. We need to use the analogue of Lemma 2.2.5 for $\mathrm{End}_{\mathbf{H}^{\mathrm{fr}}(k)^{\mathrm{sp}}}(\mathbf{1}_k)$, which holds because there is a canonical isomorphism

$$\pi_0 \mathrm{End}_{\mathbf{H}^{\mathrm{fr}}(k)^{\mathrm{sp}}}(\mathbf{1}_k) \rightarrow \pi_0 \mathrm{End}_{\mathbf{SH}(k)}(\mathbf{1}_k)$$

by [EHK⁺18b, Theorem 3.5.17].

It remains now to show that f^* is essentially surjective. Since any smooth irreducible k_{perf} -scheme is, up to a universal homeomorphism, the base change of a smooth irreducible k -scheme [Sus17, Lemma 1.12], it will suffice to show the following claim: for any universal homeomorphism of smooth schemes over k_{perf} , the induced map in $\mathbf{H}^{\mathrm{fr}}(k_{\mathrm{perf}})^{\mathrm{sp}}_{[\frac{1}{p}]}$ is invertible. By [EHK⁺18b, Theorem 3.5.13(i)] it suffices to show that the induced map in $\mathbf{SH}(k_{\mathrm{perf}})_{[\frac{1}{p}]}$ is invertible. This follows directly from Theorem 2.1.1. \square

Proof of Theorem 3.2.7. Note that we need only prove the claim when $p > 1$, and that the second claim follows from the first by stabilization. The equivalence of Corollary 2.1.4 restricts to an equivalence

$$\mathbf{SH}^{\mathrm{veff}}(S)_{[\frac{1}{p}]} \rightarrow \mathbf{SH}^{\mathrm{veff}}(S_{\mathrm{perf}})_{[\frac{1}{p}]}$$

by construction. In view of [EHK⁺18b, Theorem 3.5.13(i)], the claim follows from Lemma 3.2.9. \square

REFERENCES

- [AGV72] M. Artin, A. Grothendieck, and J.-L. Verdier, *Théorie des Topos et Cohomologie Étale des Schémas*, Lecture Notes in Mathematics, vol. 269, Springer, 1972
- [Ayo08] J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique, I*, *Astérisque* **315** (2008)
- [Bac18] T. Bachmann, *On the conservativity of the functor assigning to a motivic spectrum its motive*, *Duke Math. J.* **167** (2018), no. 8, pp. 1525–1571, <https://doi.org/10.1215/00127094-2018-0002>
- [BD15] M. Bondarko and F. Déglise, *Dimensional homotopy t-structure in motivic homotopy theory*, 2015, [arXiv:1512.06044](https://arxiv.org/abs/1512.06044)
- [BH18] T. Bachmann and M. Hoyois, *Norms in motivic homotopy theory*, 2018, [arXiv:1711.03061v4](https://arxiv.org/abs/1711.03061v4)
- [BS17] B. Bhatt and P. Scholze, *Projectivity of the Witt vector affine Grassmannian*, *Invent. Math.* **209** (2017), no. 2, pp. 329–423, <https://doi.org/10.1007/s00222-016-0710-4>
- [CD12] D.-C. Cisinski and F. Déglise, *Triangulated categories of mixed motives*, 2012, [arXiv:0912.2110v3](https://arxiv.org/abs/0912.2110v3)
- [CD15] ———, *Integral mixed motives in equal characteristic*, *Doc. Math., Extra Volume: Alexander S. Merkurjev’s Sixtieth Birthday* (2015), pp. 145–194
- [Cis13] D.-C. Cisinski, *Descente par éclatements en K-théorie invariante par homotopie*, *Ann. Math.* **177** (2013), no. 2, pp. 425–448
- [DJK18] F. Déglise, F. Jin, and A. A. Khan, *Fundamental classes in motivic homotopy theory*, 2018, [arXiv:1805.05920](https://arxiv.org/abs/1805.05920)
- [EHK⁺18a] E. Elmanto, M. Hoyois, A. A. Khan, V. Sosnilo, and M. Yakerson, *Framed transfers and motivic fundamental classes*, preprint [1809.10666](https://arxiv.org/abs/1809.10666)
- [EHK⁺18b] ———, *Motivic infinite loop spaces*, 2018, [arXiv:1711.05248v4](https://arxiv.org/abs/1711.05248v4)
- [GR71] A. Grothendieck and M. Raynaud, *Revêtements étales et groupe fondamental*, *Lecture Notes in Mathematics*, vol. 223, Springer, 1971
- [Gro67] A. Grothendieck, *Éléments de Géométrie Algébrique: IV. Étude locale des schémas et des morphismes de schémas, Quatrième partie*, *Publ. Math. I.H.É.S.* **32** (1967)
- [Gro77] ———, *Cohomologie l-adique et fonctions L*, *Lecture Notes in Mathematics*, vol. 589, Springer, 1977
- [Hoy14] M. Hoyois, *A quadratic refinement of the Grothendieck–Lefschetz–Verdier trace formula*, *Algebr. Geom. Topol.* **14** (2014), no. 6, pp. 3603–3658

- [Kha16] A. A. Khan, *Motivic homotopy theory in derived algebraic geometry*, Ph.D. thesis, Universität Duisburg-Essen, 2016, available at <https://www.preschema.com/thesis/thesis.pdf>
- [KK82] M. Knebusch and M. Kolster, *Witt rings*, Aspects of Mathematics, vol. 2, Friedr. Vieweg & Sohn, Braunschweig, 1982
- [KM18] S. Kelly and M. Morrow, *K-theory of valuation rings*, preprint 1810.12203
- [LYZ13] M. Levine, Y. Yang, and G. Zhao, *Algebraic elliptic cohomology theory and flops, I*, 2013, arXiv:1311.2159
- [MH73] J. W. Milnor and D. Husemoller, *Symmetric bilinear forms*, vol. 60, Springer, 1973
- [Mor04] F. Morel, *On the motivic π_0 of the sphere spectrum*, Axiomatic, enriched and motivic homotopy theory, NATO Sci. Ser. II Math. Phys. Chem., vol. 131, Kluwer Acad. Publ., Dordrecht, 2004, pp. 219–260, https://doi.org/10.1007/978-94-007-0948-5_7
- [Mor12] ———, *\mathbf{A}^1 -Algebraic Topology over a Field*, Lecture Notes in Mathematics, vol. 2052, Springer, 2012
- [MV99] F. Morel and V. Voevodsky, *\mathbf{A}^1 -homotopy theory of schemes*, Publ. Math. I.H.É.S. **90** (1999), pp. 45–143, preprint K-theory:0305
- [Sus17] A. Suslin, *Motivic complexes over nonperfect fields*, Ann. K-Theory **2** (2017), no. 2, pp. 277–302, <https://doi.org/10.2140/akt.2017.2.277>
- [TT90] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, The Grothendieck Festschrift III, Progress in Mathematics, vol. 88, Birkhäuser, 1990, pp. 247–435

KØBENHAVNS UNIVERSITET, INSTITUT FOR MATEMATISKE FAG, UNIVERSITETSPARKEN 5 2100 KØBENHAVN, DENMARK

E-mail address: elmanto@math.ku.dk

URL: <https://www.eldenelmanto.com/>

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, UNIVERSITÄTSSTR. 31, 93040 REGENSBURG, GERMANY

E-mail address: adeel.khan@mathematik.uni-regensburg.de

URL: <https://www.preschema.com>