

ABSOLUTE POINCARÉ DUALITY IN ÉTALE COHOMOLOGY

ADEEL A. KHAN

ABSTRACT. We extend Poincaré duality in étale cohomology from smooth schemes to regular ones. This is achieved via a formalism of trace maps for local complete intersection morphisms.

Let S be a base scheme and let Λ denote the constant sheaf $\mathbf{Z}/n\mathbf{Z}$ for an integer n which is invertible on S . For a locally of finite type¹ S -scheme X , define the Borel–Moore homology² of X (relative to S) as cohomology with coefficients in $K_X := f^!(\Lambda)$, i.e.,

$$H_*(X/S, \Lambda) = H^{-*}(X, K_X)$$

where $f : X \rightarrow S$ is the structural morphism. Our starting point is the following classical result:

Theorem 1 (Poincaré duality). *Let X be a smooth S -scheme of relative dimension d . Then there is a canonical isomorphism*

$$f^!(\Lambda) \simeq \Lambda(d)[2d]$$

in the derived category $\mathbf{D}(X_{\text{ét}}, \Lambda)$ of étale sheaves of Λ -modules on X , where $f : X \rightarrow S$ is the structural morphism. In particular, there is a canonical isomorphism

$$H_*(X/S, \Lambda) \simeq H^{2d-*}(X, \Lambda(d)). \tag{1}$$

See [SGA4, Exp. XVIII, Thm. 3.2.5], and [LZ, Thm. 0.1.4] in case X is not separated of finite type. Theorem 1 is proven by constructing a formalism of traces³

$$\text{tr}_f : f_!\Lambda(d)[2d] \rightarrow \Lambda$$

for flat morphisms f whose geometric fibres are of dimension $\leq d$ (see [SGA4, Exp. XVIII, Thm. 2.9]). By adjunction the trace gives rise to a *fundamental class*

$$[X] \in H_{2d}(X/S, \Lambda)(-d),$$

Date: 2021-07-19.

¹Classically, the $!$ -operations were constructed for compactifiable morphisms between quasi-compact quasi-separated schemes (see [SGA4, Exp. XVII]), which by Nagata is the class of separated morphisms of finite type. In [LZ], they were extended to locally of finite type morphisms.

²If S is regular and $f : X \rightarrow S$ is separated of finite type, then K_X is a dualizing complex on X by [ILO, Exp. XVII, Thm. 0.2]. Professor Illusie has informed me that the definition of Borel–Moore homology as cohomology with coefficients in K_X is in fact due to Grothendieck. See also [La1, §2].

³Throughout the note, all functors are implicitly derived (wherever necessary).

and the isomorphism (1) can be realized as cap product with $[X]$.

In this paper our goal is to prove an “absolute” version of Theorem 1, where X is a regular scheme over a fixed regular base scheme S , and the morphism $f : X \rightarrow S$ is only assumed to be locally of finite type. To that end we construct a formalism of traces for *local complete intersection* morphisms.

Theorem A. *Let X and Y be schemes on which n is invertible. For every local complete intersection morphism $f : X \rightarrow Y$ of relative virtual dimension d , there is a canonical morphism*

$$\mathrm{tr}_f : f_! \Lambda(d)[2d] \rightarrow \Lambda,$$

in $\mathbf{D}(Y_{\text{ét}}, \Lambda)$ satisfying the following properties:

- (i) *Functoriality. Given another local complete intersection morphism $g : Y \rightarrow Z$ of relative virtual dimension e , the composite $g \circ f$ is a local complete intersection morphism of relative virtual dimension $d + e$, and there is a commutative diagram*

$$\begin{array}{ccc} g_! f_! \Lambda(d)[2d](e)[2e] & \longrightarrow & g_! \Lambda(e)[2e] \\ \parallel & & \downarrow \mathrm{tr}_g \\ (g \circ f)_! \Lambda(d+e)[2d+2e] & \xrightarrow{\mathrm{tr}_{g \circ f}} & \Lambda \end{array}$$

in $\mathbf{D}(Z_{\text{ét}}, \Lambda)$. If $f = \mathrm{id}_X$, then $\mathrm{tr}_f = \mathrm{id} : \Lambda \rightarrow \Lambda$.

- (ii) *Transverse base change. Given any morphism $q : Y' \rightarrow Y$, form the cartesian square*

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ \downarrow p & & \downarrow q \\ X & \xrightarrow{f} & Y. \end{array}$$

If this square is Tor-independent (e.g. if q is flat), then g is a local complete intersection morphism of relative virtual dimension d , and there is a commutative square

$$\begin{array}{ccc} g_! \Lambda(d)[2d] & \xrightarrow{\mathrm{tr}_g} & \Lambda \\ \parallel & & \parallel \\ q^* f_! \Lambda(d)[2d] & \xrightarrow{q^*(\mathrm{tr}_f)} & q^* \Lambda \end{array}$$

in $\mathbf{D}(Y'_{\text{ét}}, \Lambda)$.

- (iii) *Purity. Denote by*

$$\mathrm{gys}_f : \Lambda(d)[2d] \rightarrow f^! \Lambda$$

the morphism in $\mathbf{D}(Y_{\text{ét}}, \Lambda)$ obtained from tr_f by transposition. If f is smooth, or if X and Y are regular, then gys_f is an isomorphism.

- (iv) *If f is smooth, then tr_f agrees with the trace morphism of [SGA4, Exp. XVIII, Thm. 2.9] (or rather [LZ, Thm. 0.1.4] in the non-compactifiable case).*

- (v) If f is a regular closed immersion, then gys_f coincides with the Gysin morphism $\text{Cl}_f : \Lambda \rightarrow f^! \Lambda(-d)[-2d]$ constructed in [ILO, §2.3] and [Azu, §1] (that is, $\text{gys}_f = \text{Cl}_f(d)[2d]$). In particular, it refines the local cycle class of [Cycle, §2.2].

Here local complete intersection (lci) morphisms are defined as in [SGA6, Exp. VIII, §1, Déf. 1.1]. For us the relevant description will be as follows: a morphism of schemes is lci if and only if it is locally of finite presentation and has perfect relative cotangent complex of Tor-amplitude $[-1, 0]$ (under cohomological grading conventions). See e.g. [KRy, Prop. 2.3.14] for this equivalence.⁴

Remark 1. Fix a base scheme S on which n is invertible. From Theorem A we can now read off:

- (i) If X is an lci S -scheme of relative virtual dimension d then it admits a *fundamental class*

$$[X] \in \mathbb{H}_{2d}(X/S, \Lambda)(-d),$$

given by the morphism $\text{gys}_p(-d)[-2d] : \Lambda \rightarrow p^! \Lambda(-d)[-2d]$, where $p : X \rightarrow S$ is the structural morphism.⁵

- (ii) If X and S are regular and $d = \dim(X) - \dim(S)$, then $\text{gys}_p : p^! \Lambda \simeq \Lambda(d)[2d]$ gives rise to canonical isomorphisms (“absolute Poincaré duality”)

$$\cap[X] : \mathbb{H}^*(X, \Lambda) \rightarrow \mathbb{H}_{2d-*}(X/S, \Lambda)(-d).$$

- (iii) For any lci morphism $f : X \rightarrow Y$ between S -schemes, $\text{gys}_f : \Lambda(d)[2d] \rightarrow f^! \Lambda$ gives rise to Gysin pull-backs

$$f^! : \mathbb{H}_*(Y/S, \Lambda) \rightarrow \mathbb{H}_{*+2d}(X/S, \Lambda)(-d).$$

- (iv) For any *proper* lci morphism $f : X \rightarrow Y$ of relative virtual dimension d , tr_f gives rise to Gysin push-forwards in cohomology

$$f_! : \mathbb{H}^*(X, \Lambda) \rightarrow \mathbb{H}^{*-2d}(Y, \Lambda(-d)).$$

1. DEFORMATION TO THE NORMAL STACK

The main ingredient is *deformation to the normal stack*, a variant of deformation to the normal cone that makes sense not just for closed immersions.

Given an lci morphism $f : X \rightarrow Y$ of schemes, the normal stack $N_{X/Y}$ is the “total space” of the (-1) -shifted cotangent complex $L_{X/Y}[-1]$. To make sense of this, recall that the total space construction $\mathcal{E} \mapsto \mathbf{V}_X(\mathcal{E}) = \text{Spec}_X(\text{Sym}_{\mathcal{O}_X}(\mathcal{E}))$ defines an equivalence between finite locally free sheaves and vector bundles over X . This extends to an equivalence between perfect

⁴Surprisingly, I do not know an earlier reference. In fact, it seems to be a common misconception that this requires noetherianness or that it is related to Quillen’s conjecture.

⁵If S is a field and X is quasi-projective, then this is the image of the fundamental class in the Chow group $A_d(X)$ by the cycle class map.

complexes of Tor-amplitude $[0, 1]$ and vector bundle stacks over X , so that we can write

$$N_{X/Y} := \mathbf{V}_X(L_{X/Y}[-1]).$$

See [Kh, §1.3], [BF, §2], [SGA4, Exp. XVIII, §1.4]. In [BF] this is called the “intrinsic normal cone” or “intrinsic normal sheaf” (they agree for lci morphisms).

If f is a closed immersion, then $L_{X/Y}[-1]$ is just the conormal sheaf in degree zero so $N_{X/Y}$ is just the normal bundle. In general, $L_{X/Y}[-1]$ will typically have nonzero cohomology in degree 1, which is why $N_{X/Y}$ will only exist as an algebraic stack. For example if f is smooth then $L_{X/Y}[-1]$ is the cotangent sheaf in degree -1 , so $BT_{X/Y}$ is the classifying stack of the tangent bundle. If there is a global factorization of f through a regular immersion $i : X \hookrightarrow M$ and a smooth morphism $p : M \rightarrow Y$, then $N_{X/Y}$ is isomorphic to the stack quotient

$$N_{X/Y} \simeq [N_{X/M}/i^*T_{M/Y}]$$

where $N_{X/M}$ is the normal bundle of i and $T_{M/Y}$ is the relative tangent bundle of p . No choices are involved in the definitions of $L_{X/Y}$ and $N_{X/Y}$, i.e., they are intrinsic to f .

Deformation to the normal stack is an \mathbf{A}^1 -family of algebraic stacks which deforms $f : X \rightarrow Y$ to the zero section $0 : X \rightarrow N_{X/Y}$. That is, it is a commutative diagram of algebraic stacks

$$\begin{array}{ccccc} X & \xrightarrow{0} & X \times \mathbf{A}^1 & \longleftarrow & X \times \mathbf{G}_m \\ \downarrow 0 & & \downarrow & & \downarrow f \times \text{id} \\ N_{X/Y} & \xrightarrow{\hat{i}} & D_{X/Y} & \xleftarrow{\hat{j}} & Y \times \mathbf{G}_m \\ \downarrow & & \downarrow & & \parallel \\ Y & \xrightarrow{0} & Y \times \mathbf{A}^1 & \longleftarrow & Y \times \mathbf{G}_m \end{array} \quad (2)$$

where each square is cartesian (and Tor-independent). If f is a closed immersion then this is Verdier’s deformation to the normal cone. See [Kh, §1.4], [Kr, §5.1], [Ma, Thm. 2.31].

2. BOREL–MOORE HOMOLOGY OF STACKS

Using the extension of the six operations to algebraic stacks defined in [LZ]⁶ we can define Borel–Moore homology of an algebraic stack \mathcal{X} (locally of finite type over some base S) again by the formula

$$H_k(\mathcal{X}/S, \Lambda) = H^{-k}(\mathcal{X}, p^!(\Lambda))$$

where $p : \mathcal{X} \rightarrow S$ is the structural morphism. Equivalently, these are the homology groups of the complexes

$$\mathbf{R}\Gamma(\mathcal{X}/S, \Lambda) := \mathbf{R}\Gamma(\mathcal{X}, p^!(\Lambda)) \simeq \mathbf{R}\varprojlim_{(T,t)} \mathbf{R}\Gamma(T, (p \circ t)^!(\Lambda))(-d_t)[-2d_t],$$

⁶If we work over a base satisfying some strong hypotheses, which hold e.g. for spectra of finite or separably closed fields, then we can also use the formalism of [LO]; cf. [LZ, §6.5].

where the right-hand side is the homotopy limit over pairs (T, t) where T is a scheme and $t: T \rightarrow \mathcal{X}$ is a smooth morphism of relative dimension d_t . (By Zariski descent, T can also be taken affine.) Note that this second description only requires an enhancement of the assignment $X \mapsto \mathbf{D}(X_{\text{ét}}, \Lambda)$, $f \mapsto f^!$ to a presheaf of ∞ -categories on the category of schemes and locally of finite type morphisms. Aside from [LZ], there are also various other approaches to such a construction, see e.g. [GR, AGV].

It is straightforward to deduce that the localization exact triangle extends to stacks:

Proposition 1 (Localization). *If $i: \mathcal{Z} \rightarrow \mathcal{X}$ is a closed immersion with open complement $j: \mathcal{U} \rightarrow \mathcal{X}$, then we have an exact triangle*

$$\mathbf{R}\Gamma(\mathcal{Z}/S, \Lambda) \xrightarrow{i_*} \mathbf{R}\Gamma(\mathcal{X}/S, \Lambda) \xrightarrow{j^!} \mathbf{R}\Gamma(\mathcal{U}/S, \Lambda),$$

whence a long exact sequence

$$\cdots \rightarrow \mathbf{H}_k(\mathcal{Z}/S, \Lambda) \xrightarrow{i_*} \mathbf{H}_k(\mathcal{X}/S, \Lambda) \xrightarrow{j^!} \mathbf{H}_k(\mathcal{U}/S, \Lambda) \xrightarrow{\partial} \mathbf{H}_{k-1}(\mathcal{Z}/S, \Lambda) \rightarrow \cdots.$$

For example, consider the closed/open pair (\hat{i}, \hat{j}) from (2). The boundary map in this case gives rise to a specialization map

$$\begin{aligned} \text{sp}_{X/Y} : \mathbf{H}_k(Y/Y, \Lambda) &\xrightarrow{\text{incl}} \mathbf{H}_k(Y/Y, \Lambda) \oplus \mathbf{H}_{k-1}(Y/Y, \Lambda)(1) \\ &\simeq \mathbf{H}_{k+1}(Y \times \mathbf{G}_m/Y, \Lambda) \xrightarrow{\partial} \mathbf{H}_k(N_{X/Y}/Y, \Lambda). \end{aligned} \quad (3)$$

We will also need homotopy invariance for vector bundle stacks:

Proposition 2. *Let \mathcal{E} be the total space of a perfect complex of Tor-amplitude $[0, 1]$ over X , say of virtual rank r . Then there is a canonical isomorphism in the derived category of Λ -modules*

$$\mathbf{R}\Gamma(\mathcal{E}/Y, \Lambda) \simeq \mathbf{R}\Gamma(X/Y, \Lambda)(r)[2r].$$

In particular,

$$\mathbf{H}_k(\mathcal{E}/Y, \Lambda) \simeq \mathbf{H}_{k-2r}(X/Y, \Lambda)(r)$$

for all $k \in \mathbf{Z}$.

Proof. Since the projection $\pi: \mathcal{E} \rightarrow X$ is smooth of relative dimension r , we have the Poincaré duality isomorphism

$$\mathbf{R}\Gamma(\mathcal{E}, \pi^! f^! \Lambda) \simeq \mathbf{R}\Gamma(\mathcal{E}, f^! \Lambda)(r)[2r],$$

where there is an implicit π^* on the right-hand side. This is the homotopy limit over (T, t) of the Poincaré duality isomorphisms

$$\mathbf{R}\Gamma(T, (\pi \circ t)^! f^! \Lambda)(-d_t)[-2d_t] \simeq \mathbf{R}\Gamma(T, f^! \Lambda)(r)[2r]$$

for the smooth morphism $\pi \circ t: T \rightarrow \mathcal{X} \rightarrow Y$ of relative dimension $d_t + r$.⁷

⁷Note that to form this homotopy limit, we need the Poincaré duality isomorphism for schemes to be functorial in a homotopy coherent sense; however, this coherence comes for free using a standard t-structure argument, see [LZ, Thm. 6.2.9, Rem. 4.1.10].

Secondly, there is a canonical map

$$\pi^* : \mathbf{R}\Gamma(X, f^!\Lambda) \rightarrow \mathbf{R}\Gamma(\mathcal{E}, f^!\Lambda)$$

which (as a consequence of étale descent) can be described as the homotopy limit of the maps π_U^* , where $\pi_U : \mathcal{E} \times_Y U \rightarrow U$, taken over smooth morphisms $U \rightarrow Y$ with U affine. Therefore the claim is local on Y and we may assume that the perfect complex defining \mathcal{E} admits a global resolution, so that \mathcal{E} is globally the stack quotient $[E^1/E^0]$ of a vector bundle morphism $E^0 \rightarrow E^1$. In this case π^* factors through isomorphisms

$$\mathbf{R}\Gamma(X, f^!\Lambda) \rightarrow \mathbf{R}\Gamma(E^1, f^!\Lambda) \leftarrow \mathbf{R}\Gamma(\mathcal{E}, f^!\Lambda)$$

by homotopy invariance for the vector bundle $E^1 \rightarrow X$ (follows by descent from the case of trivial bundles, see [SGA4, Exp. XV, Cor. 2.2]) and for the E^0 -torsor $E^1 \rightarrow \mathcal{E}$ (can be checked after base change to affines, over which vector bundle torsors are split). \square

3. THE CONSTRUCTION

We return to the situation of an lci morphism $f : X \rightarrow Y$, say of relative virtual dimension d . The (-1) -shifted cotangent complex $L_{X/Y}[-1]$ is perfect of Tor-amplitude $[0, 1]$ (of virtual rank $-d$), so Proposition 2 yields a canonical isomorphism

$$\mathbf{H}_k(N_{X/Y}/Y, \Lambda) \simeq \mathbf{H}_{k+2d}(X/Y, \Lambda)(-d).$$

Combining this with the specialization map (3) produces now a canonical map

$$\mathbf{H}_k(Y/Y, \Lambda) \xrightarrow{\text{sp}_{X/Y}} \mathbf{H}_k(N_{X/Y}/Y, \Lambda) \simeq \mathbf{H}_{k+2d}(X/Y, \Lambda)(-d).$$

In particular, the image of the unit $1 \in \mathbf{H}_0(Y/Y, \Lambda)$ gives rise to a canonical element (a relative fundamental class)

$$[X/Y] \in \mathbf{H}_{2d}(X/Y, \Lambda)(-d). \quad (4)$$

Our Gysin morphism is then the corresponding morphism

$$\text{gys}_f : \Lambda(d)[2d] \rightarrow f^!\Lambda \quad (5)$$

in $\mathbf{D}(X_{\text{ét}}, \Lambda)$, and the trace morphism $\text{tr}_f : f_!\Lambda(d)[2d] \rightarrow \Lambda$ is its transpose.

It will also be useful to note that these can be refined to natural transformations

$$\text{gys}_f : f^*(d)[2d] \rightarrow f^! \quad (6)$$

$$\text{tr}_f : f_! f^*(d)[2d] \rightarrow \text{id}. \quad (7)$$

For example, tr_f is the composite

$$f_! f^*(-)(d)[2d] \simeq (-) \otimes f_!\Lambda(d)[2d] \xrightarrow{\text{id} \otimes \text{tr}_f} (-) \otimes \Lambda = \text{id}$$

where the isomorphism is the projection formula. Note that when (5) is invertible, (6) will also be invertible on dualizable objects in $\mathbf{D}(Y_{\text{ét}}, \Lambda)$ (but not necessarily on arbitrary ones).

4. PROOFS OF THE ASSERTED PROPERTIES

We begin by noting that, in case f is a closed immersion, our construction of the Gysin morphism obviously coincides with that of [DJK, §3.2], which itself agrees with Gabber's construction [ILO, Exp. XVI, §2.3] by [DJK, 4.4.3]. The base change and functoriality properties are proven exactly as in the case of closed immersions, using respectively Tor-independent base change of the deformation space $D_{X/Y}$ (see [Kh, Thm. 1.3(ii)]) and the double deformation space associated to lci morphisms $X \rightarrow Y \rightarrow Z$,

$$D_{X/Y/Z} := D_{D_{X/Z} \times_Z Y / D_{X/Z}},$$

the deformation to the normal stack of the morphism $D_{X/Z} \times_Z Y \rightarrow D_{X/Z}$. See the proof of [DJK, Thm. 3.2.21].

Let us show that if f is smooth of relative dimension d , then gys_f is the Poincaré duality isomorphism $f^!(\Lambda) \simeq \Lambda(d)[2d]$. Form the cartesian square

$$\begin{array}{ccc} X \times_Y X & \xrightarrow{\text{pr}_2} & X \\ \downarrow \text{pr}_1 & & \downarrow f \\ X & \xrightarrow{f} & Y \end{array}$$

The diagonal morphism $\Delta : X \rightarrow X \times_Y X$ is lci of relative virtual dimension $-d$ and the natural transformation $\text{tr}_\Delta : \Delta_! \Delta^*(-d)[-2d] \rightarrow \text{id}$ (7) gives rise to

$$\eta_f : \text{id} = \text{pr}_{2,!} \Delta_! \Delta^* \text{pr}_1^* \xrightarrow{\text{tr}_\Delta} \text{pr}_{2,!} \text{pr}_1^*(d)[2d] \simeq f^* f_!(d)[2d].$$

We claim that η_f and tr_f form the unit and counit of an adjunction $(f_!, f^*(d)[2d])$. Indeed, it is easy to check that both composites

$$\begin{array}{ccccc} f_! & \xrightarrow{f_!(\eta_f)} & f_! f^* f_!(d)[2d] & \xrightarrow{\text{tr}_f * f_!} & f_! \\ f^*(d)[2d] & \xrightarrow{\eta_f * f^*(d)[2d]} & f^* f_! f^*(2d)[4d] & \xrightarrow{f^* * \text{tr}_f(d)[2d]} & f^*(d)[2d] \end{array}$$

are identity by using the functoriality of the trace for the composite $\text{pr}_1 \circ \Delta$ (resp. for the composite $\text{pr}_2 \circ \Delta$) and by base change for the trace of f . This argument shows not only that gys_f is an isomorphism but also that it agrees with the Poincaré duality isomorphisms of [SGA4, Exp. XVIII, Thm. 3.2.5] and [LZ, Thm. 0.1.4], or equivalently that tr_f agrees with the trace of [SGA4, Exp. XVIII, Thm. 2.9] or [LZ, Thm. 0.1.4]: indeed, both are counits for the same adjunction.

It remains to show that if X and Y are regular (in which case $f : X \rightarrow Y$ is automatically lci), then gys_f gives the isomorphism $f^! \Lambda \simeq \Lambda(d)[2d]$ asserted in Theorem A(iii). But invertibility of gys_f can be checked after inverse image along a Zariski cover, and by functoriality we have for any open

immersion $j : U \hookrightarrow X$ a commutative diagram

$$\begin{array}{ccccc} \Lambda(d)[2d] & \xrightarrow{\text{gys}_j} & j^! \Lambda(d)[2d] & \xrightarrow{j^!(\text{gys}_f)} & j^! f^! \Lambda \\ \parallel & & & & \parallel \\ \Lambda(d)[2d] & \xrightarrow{\text{gys}_{f \circ j}} & & & (f \circ j)^! \Lambda. \end{array}$$

where gys_j is invertible. Thus we may localize on X and choose a global factorization through a closed immersion $i : X \hookrightarrow X'$ and a smooth morphism $p : X' \rightarrow Y$. By functoriality of Gysin morphisms again and the fact that gys_p is an isomorphism by above, we reduce to the case of a closed immersion between regular schemes (note that X' is still regular). Finally, since gys_f agrees with Gabber's construction in this case, the claim now follows from absolute purity [ILO, Exp. XVI, Thm. 3.1.1].

5. REMARKS

Using the formalism of [LZ], our construction of the traces tr_f immediately extends to the case where the schemes X and Y are algebraic stacks. Absolute Poincaré duality also extends to regular algebraic stacks with the same proof.

We can also allow X and Y to be derived (schemes or stacks), and $f : X \rightarrow Y$ to be any *quasi-smooth* morphism. Indeed, an lci morphism is precisely a quasi-smooth morphism whose source and target happen to be classical (underived). The construction of Theorem A goes through mutatis mutandis, since the deformation space $D_{X/Y}$ exists in that setting (see [Kh, §1.4]): it is simply the Weil restriction of X along $0 : Y \hookrightarrow Y \times \mathbf{A}^1$ (in the derived sense). For a quasi-smooth morphism, the trace is a kind of categorification of Kontsevich's virtual fundamental class (cf. (4)) and gives rise for example to the Gromov–Witten theory of smooth projective varieties in arbitrary characteristic. On the other hand, absolute Poincaré duality does not hold for derived schemes whose classical truncations are not regular.

Finally, the construction can be refined from étale cohomology to motivic cohomology. For this one can use the limit-extended motivic cohomology of algebraic stacks defined in [KRa, §12] as a substitute for [LZ]. Note that the trace formalism for *flat* maps (as developed in [SGA4, Exp. XVIII, Thm. 2.9]) has recently been extended to motivic cohomology by Abe (see [Abe]). Note that in this setting our proof of absolute Poincaré duality goes through only in equicharacteristic, since absolute purity in motivic cohomology is open in general (see [DFJK, Thm. C.1] for the equicharacteristic case).

In the setting of rational and étale motivic cohomology, the results of this paper appeared in a somewhat different form in the preprint [Kh]. While rewriting the latter, I felt that the absolute Poincaré duality result was fairly disjoint from the main goals of [Kh]. The present paper is an attempt to give a short and self-contained account without using the language of motives or derived algebraic geometry.

In a future paper, I will explain how to use deformation to the normal stack to generalize Verdier’s specialization functor [Ve]. This will be combined with a derived version of Laumon’s homogeneous Fourier transform [La2] to give an analogue of microlocalization in the sense of Kashiwara–Schapira [KS] for singular schemes.

6. ACKNOWLEDGMENTS

I would like to thank Luc Illusie and Joël Riou for helpful comments on a draft.

These results were first announced during a conference at the Euler International Mathematical Institute in St. Petersburg, Russia in 2019. I would like to thank the organizers for the invitation to speak there. At the time, I was partially supported by SFB 1085 Higher Invariants, Universität Regensburg.

REFERENCES

- [AGV] J. Ayoub, M. Gallauer, A. Vezzani, *The six-functor formalism for rigid analytic motives*. [arXiv:2010.15004](https://arxiv.org/abs/2010.15004) (2020).
- [Abe] T. Abe, *Trace formalism for motivic cohomology*.
- [Azu] K. Fujiwara, *A proof of the absolute purity conjecture (after Gabber)*, in: Algebraic geometry 2000, Azumino. Adv. Stud. Pure Math. **36** (2002), 153–183.
- [BF] K. Behrend, B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128** (1997), no. 1, 45–88.
- [Cycle] A. Grothendieck, *La classe de cohomologie associée à un cycle (rédigé par P. Deligne)*, in: Cohomologie étale. Séminaire de Géométrie Algébrique du Bois-Marie (SGA 4 $\frac{1}{2}$). Lecture Notes in Mathematics **569**, Springer (1977).
- [DFJK] F. Déglise, J. Fasel, F. Jin, A. A. Khan, *On the rational motivic homotopy category*. J. Éc. Polytech., Math. **8** (2021), 533–583.
- [DJK] F. Déglise, F. Jin, A. A. Khan, *Fundamental classes in motivic homotopy theory*. J. Eur. Math. Soc. (JEMS), to appear. [arXiv:1805.05920](https://arxiv.org/abs/1805.05920) (2018).
- [GR] D. Gaißgory, N. Rozenblyum, *A study in derived algebraic geometry. Vol. I. Correspondences and duality*. Math. Surv. Mono. **221** (2017).
- [ILO] L. Illusie, Y. Laszlo, F. Orgogozo (eds.), *Travaux de Gabber sur l’uniformisation locale et la cohomologie étale des schémas quasi-excellents*, Séminaire à l’École Polytechnique 2006–2008. Astérisque **363–364**, Soc. Math. France (2014).
- [KS] M. Kashiwara, P. Schapira, *Sheaves on manifolds* (1990).
- [KRa] A. A. Khan, C. Ravi, *Generalized cohomology theories for algebraic stacks*. [arXiv:2106.15001](https://arxiv.org/abs/2106.15001) (2021).
- [KRy] A. A. Khan, D. Rydh, *Virtual Cartier divisors and blow-ups*. [arXiv:1802.05702](https://arxiv.org/abs/1802.05702) (2018).
- [Kh] A. A. Khan, *Virtual fundamental classes for derived stacks I*. [arXiv:1909.01332](https://arxiv.org/abs/1909.01332) (2019).
- [Kr] A. Kresch, *Cycle groups for Artin stacks*. Invent. Math. **138** (1999), no. 3, 495–536.
- [LO] Y. Laszlo, M. Olsson, *The six operations for sheaves on Artin stacks. I. Finite coefficients*. Publ. Math. Inst. Hautes Études Sci., no. **107** (2008), 109–168.
- [LZ] Y. Liu, W. Zheng, *Enhanced six operations and base change theorem for higher Artin stacks*. [arXiv:1211.5948](https://arxiv.org/abs/1211.5948) (2012).
- [La1] G. Laumon, *Homologie étale*, Exposé VIII in Séminaire de géométrie analytique (Ecole Norm. Sup., Paris, 1974–75), Astérisque **36–37**, 163–188. Soc. Math. France (1976).

- [La2] G. Laumon, *Transformation de Fourier homogène*. Bull. Soc. Math. Fr. **131** (2003), no. 4, 527–551.
- [Ma] C. Manolache, *Virtual pull-backs*. J. Algebraic Geom. **21** (2012), no. 2, 201–245.
- [SGA4] M. Artin, A. Grothendieck, J. L. Verdier (eds.), *Théorie des topos et cohomologie étale des schemas* (SGA 4). Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964. Lecture Notes in Mathematics **269, 270, 305**, Springer (1972).
- [SGA5] A. Grothendieck (ed.), *Cohomologie l -adique et fonctions L* (SGA 5). Séminaire de Géométrie Algébrique du Bois-Marie 1965–1966. Lecture Notes in Mathematics **589**, Springer (1977).
- [SGA6] P. Berthelot, A. Grothendieck, L. Illusie (eds.), *Théorie des intersections and théorème de Riemann–Roch*. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6). Lecture Notes in Mathematics **225**, Springer (1971).
- [Ve] J.-L. Verdier, *Spécialisation de faisceaux et monodromie modérée*. Astérisque **101-102** (1983), 332–364.

Institute of Mathematics, Academia Sinica, Taipei 10617, Taiwan