

K-THEORY AND G-THEORY OF DERIVED ALGEBRAIC STACKS

ADEEL A. KHAN

ABSTRACT. These are some notes on the basic properties of algebraic K-theory and G-theory of derived algebraic spaces and stacks, and the theory of fundamental classes in this setting.

Introduction	2
1. Perfect and coherent complexes	3
1.1. Complexes over derived commutative rings	3
1.2. Quasi-coherent, perfect, and coherent complexes	5
1.3. Tensor, Hom, and inverse/direct images	6
1.4. Finiteness of Tor-amplitude	8
1.5. Finiteness of cohomological dimension	8
1.6. Cohomological properness	10
1.7. The resolution property	11
1.8. Perfect stacks	12
1.9. Perfectness criteria	13
1.10. Continuity	16
2. Algebraic K-theory	18
2.1. Definitions	18
2.2. Operations	20
2.3. Projective bundles and blow-ups	21
2.4. Excising closed subsets	23
2.5. Excising open subsets	24
2.6. Bass fundamental sequence	25
2.7. Continuity	26
2.8. Nil-invariance	26
3. G-theory	27
3.1. Definition and basic properties	27
3.2. Operations	28
3.3. Excision	29
3.4. Projective bundles and blow-ups	30
3.5. Homotopy invariance	31
4. Homotopy invariance and singularities	32
4.1. Weibel's conjecture	32
4.2. Homotopy invariant K-theory	33
4.3. Properties of KH	34
5. Rational étale K-theory and G-theory	35
5.1. Descent on algebraic spaces	36
5.2. Étale K-theory and G-theory	37
5.3. Proper codescent	38
5.4. Non-representable direct image	40

6. Virtual fundamental classes	41
6.1. Fundamental classes in G-theory	41
6.2. Fundamental classes via deformation to the normal stack	42
6.3. Grothendieck–Riemann–Roch formulas	43
6.4. K-theoretic fundamental classes	45
6.5. The γ -filtration and Chow cohomology of singular schemes	46
References	49

INTRODUCTION

In [Kh5] I studied the cohomology and Borel–Moore homology of derived schemes and algebraic spaces, as well as Borel-type extensions to derived algebraic stacks. In these notes I describe the K-theoretic counterpart to that formalism. In this analogy, the K-theory of perfect complexes behaves like cohomology while G-theory (K-theory of coherent sheaves) behaves like Borel–Moore homology. This relationship can actually be made quite precise using the formalism of motivic categories: the theory of *KGL-modules* provides a category of coefficients for K-theory¹, and Borel–Moore homology in that setting (i.e., cohomology with coefficients in the dualizing complex) is G-theory. See [CD, §13.3] and [Ji].

The notes are almost entirely expository and were originally prepared as background material for the paper [Kh5]. I begin in Section 1 with some preliminaries on quasi-coherent complexes on derived stacks and various finiteness conditions. Thanks primarily to work of Hall and Rydh over the past several years, we can work with very general stacks. I also review some new compact generation results for stacks, obtained jointly with Ravi.

In the **second** and **third** sections I define K-theory and G-theory and record their basic properties. Most of these go back to [SGA6], [Q2], and [TT] in the case of classical schemes, and to [To1, To2, KrRa, HoKr] in the case of classical stacks.

The **fourth** section discusses the failure of homotopy invariance in K-theory of singular spaces and some of its ramifications. I explain how *forcing* homotopy invariance for arbitrary spaces results in a new cohomology theory, first introduced by Weibel [We2] for classical schemes, that agrees with K-theory on nonsingular spaces but behaves more “correctly” on singular ones.

In the **fifth** section I explain how K-theory and G-theory acquire some useful descent properties after passage to rational coefficients. This leads to étale-local variants of K-theory and G-theory of stacks, which do not agree however with K-theory and G-theory even rationally. A subtlety here is the existence

¹or rather its homotopy invariant version KH (see Subsect. 4.2)

and behaviour of direct images in étale G-theory, which are not compatible with those in G-theory. Note that, for stacks, these étale-localized theories are the ones that are actually compatible with the étale motivic theories studied in [Kh5].

The most important aspect of the formalism in [Kh5] is the theory of fundamental classes. The K-theoretic counterpart to that part of the story, at least some of which is known to the experts, is developed in Section 6. The K-theoretic virtual structure sheaf of [Lee] and its basic properties fall out of this formalism for free. I also give another description of Gysin maps in G-theory using deformation to the normal stack. Then I recall some variants of the Grothendieck–Riemann–Roch theorem. One formulation, proven in [Kh5], compares the virtual structure sheaf with the virtual fundamental class in Borel–Moore homology (or the Chow groups). Another is a direct generalization of the one proven in [SGA6] and involves the behaviour of K-theoretic fundamental classes with respect to the γ -filtration. This inspires some unsolicited speculations on a theory of derived algebraic cycles.

Conventions. I generally tried to follow the same conventions and notation as in [Kh5]. Since a substantial revision to that paper is in preparation, I was not always able to achieve this.

Acknowledgments. Thanks to Mauro Porta and Tony Yue Yu for many questions and suggestions about [Kh5], which eventually led to these notes being written. Thanks to Marc Hoyois, Charanya Ravi, and David Rydh for helpful conversations about stacks over the years. An additional thanks to Charanya for pointing out many typos in a previous draft and for help with the material on perfect stacks.

1. PERFECT AND COHERENT COMPLEXES

1.1. Complexes over derived commutative rings. Let A be a derived commutative ring. We write $\mathbf{D}(A)$ for the derived ∞ -category of complexes over A , defined e.g. as in [Lu4, §25.2.1], [HLP, App. A], or [TV, §2.4].

We will say a complex $M \in \mathbf{D}(A)$ is *connective*, resp. *coconnective*, if we have $\pi_i(M) = 0$ for all $i < 0$, resp. for all $i > 0$. More generally we say M is *n-connective*, resp. *n-coconnective*, if we have $\pi_i(M) = 0$ for all $i < n$, resp. for all $i > n$. We write

$$\mathbf{D}(A)_{\geq n} \subseteq \mathbf{D}(A), \quad \mathbf{D}(A)_{\leq n} \subseteq \mathbf{D}(A),$$

for the respective full subcategories. If M is both connective and coconnective, i.e., if $\pi_i(M) = 0$ for all $i \neq 0$, then we say M is *discrete*. The full subcategory of discrete complexes

$$\mathbf{D}(A)^\heartsuit \subseteq \mathbf{D}(A)$$

is equivalent to the abelian category $\text{Mod}_{\pi_0(A)}$ of $\pi_0(A)$ -modules. What we have just described is of course nothing else than the standard t-structure on $\mathbf{D}(A)$.

The cohomologically oriented reader will want to write

$$H^{-i}(M) := \pi_i(M)$$

and reset the notation as follows:

$$\mathbf{D}(A)^{\leq n} := \mathbf{D}(A)_{\geq -n}, \quad \mathbf{D}(A)^{\geq n} := \mathbf{D}(A)_{\leq -n}.$$

We recall some finiteness conditions on complexes.

Definition 1.1. A complex M over A is *perfect* if it is in the thick subcategory of $\mathbf{D}(A)$ generated by A . That is, if it is built out of A under finite (co)limits and direct summands.

To define (pseudo)coherent complexes, it is convenient (but not necessary) to assume that A is noetherian. Here this means that the ordinary commutative ring $\pi_0(A)$ is noetherian and that the homotopy groups $\pi_i(A)$ are finitely generated as $\pi_0(A)$ -modules for all $i \geq 0$. Whenever we discuss (pseudo)coherent complexes below, the reader should either assume the ring is noetherian² or should replace the definition of pseudocoherence below by [Lu3, Def. 7.2.4.10].

Definition 1.2. (i) A complex M over A is *pseudocoherent*³ if it is eventually connective, i.e., $\pi_i(M) = 0$ for $i < 0$, and $\pi_i(M)$ is finitely generated as a $\pi_0(A)$ -module for all i . (ii) A complex M over A is *coherent* if it is pseudocoherent and also eventually coconnective, i.e., $\pi_i(M) = 0$ for $i \gg 0$.

Definition 1.3. (i) A complex M over A is of *Tor-amplitude* $\leq n$ if for every discrete complex $N \in \mathbf{D}(A)^\heartsuit$, the derived tensor product $M \otimes_A^{\mathbf{L}} N$ is n -coconnective. Equivalently, for every coconnective complex $N \in \mathbf{D}(A)_{\leq 0}$, the derived tensor product $M \otimes_A^{\mathbf{L}} N$ is n -coconnective (see [Lu3, Prop. 7.2.4.23 (5)]). (ii) A complex M over A is of *finite Tor-amplitude* if it is of Tor-amplitude $\leq n$ for some n .

The following lemma summarizes the relationships between these finiteness conditions.

Lemma 1.4.

- (i) *The property of (pseudo)coherence is stable under finite (co)limits and direct summands. In other words, the (pseudo)coherent complexes form a thick subcategory of $\mathbf{D}(A)$.*
- (ii) *Every perfect complex over A is pseudocoherent.*
- (iii) *A pseudocoherent complex over A is perfect if and only if it is of finite Tor-amplitude.*
- (iv) *If A is eventually coconnective, then every perfect complex over A is coherent.*

²Noetherianness can be replaced harmlessly by the slightly weaker property of coherence in the sense of [Lu3, Def. 7.2.4.13].

³In [Lu3, Lu4, HLP], the term “almost perfect” is used instead of pseudocoherent.

- (v) *Assume A is eventually coconnective and that every coherent complex is perfect. Then A is discrete, and regular as an ordinary commutative ring.*

Proof. (i) See [Lu3, Lem. 7.2.4.11 (1-2)]. (ii) By definition of perfectness, it suffices to show that A itself is pseudocoherent over A . If A is noetherian, then this follows from the definition of pseudocoherence; see [Lu3, 7.2.4.11 (3)] for the non-noetherian case. (iii) See [Lu3, Prop. 7.2.4.23(4)]. (iv) Again it suffices to show that A is coherent over itself, which follows from the definitions when A is eventually coconnective. (v) See [Lu4, Lem. 11.3.3.3]. \square

1.2. Quasi-coherent, perfect, and coherent complexes. Let \mathcal{X} be a derived Artin stack. We write $\mathbf{D}_{\text{qc}}(\mathcal{X})$ for the stable ∞ -category of *quasi-coherent complexes* on \mathcal{X} , see e.g. [To5, Subsect. 3.1].

Over an affine derived scheme $\text{Spec}(A)$, a quasi-coherent complex \mathcal{F} is the same datum as that of the complex $\mathbf{R}\Gamma(\text{Spec}(A), \mathcal{F})$ over A , i.e., there is an equivalence

$$\mathbf{D}_{\text{qc}}(\text{Spec}(A)) \simeq \mathbf{D}(A)$$

given by $\mathcal{F} \mapsto \mathbf{R}\Gamma(\text{Spec}(A), \mathcal{F})$. We write $\mathcal{O}_X \in \mathbf{D}_{\text{qc}}(\text{Spec}(A))$ for the quasi-coherent complex whose derived global sections are $A \in \mathbf{D}(A)$.

Over a derived Artin stack \mathcal{X} , a quasi-coherent complex \mathcal{F} amounts to the data of quasi-coherent complexes $u^*(\mathcal{F}) \in \mathbf{D}_{\text{qc}}(X)$ for every smooth morphism $u : X \rightarrow \mathcal{X}$ with X affine, together with a homotopy coherent system of compatibilities between these complexes as u varies. For example, the structure sheaf $\mathcal{O}_{\mathcal{X}} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ satisfies $u^*(\mathcal{O}_{\mathcal{X}}) \simeq \mathcal{O}_X$ for every u .

Definition 1.5. (i) A quasi-coherent complex \mathcal{F} on an affine derived scheme $X = \text{Spec}(A)$ is *perfect*, *pseudocoherent*, or *coherent* if $\mathbf{R}\Gamma(X, \mathcal{F})$ has the respective property as a complex over A . (ii) A quasi-coherent complex \mathcal{F} on a derived Artin stack \mathcal{X} is *perfect*, *pseudocoherent*, or *coherent*, if $u^*(\mathcal{F})$ has the respective property for every smooth morphism $u : X \rightarrow \mathcal{X}$ from an affine X .

We write

$$\mathbf{D}_{\text{perf}}(\mathcal{X}) \subseteq \mathbf{D}_{\text{qc}}(\mathcal{X}) \text{ and } \mathbf{D}_{\text{coh}}(\mathcal{X}) \subseteq \mathbf{D}_{\text{qc}}(\mathcal{X})$$

for the full subcategories of perfect and coherent complexes on \mathcal{X} .

Remark 1.6. If \mathcal{X} is a classical Artin stack, then $\mathbf{D}_{\text{qc}}(\mathcal{X})$ can be described as the derived ∞ -category of complexes of $\mathcal{O}_{\mathcal{X}}$ -modules (on the lisse-étale topos) with quasi-coherent cohomology. See [HR2, Prop. 1.3]. In that language, $\mathbf{D}_{\text{coh}}(\mathcal{X})$ is the full subcategory of complexes with bounded and coherent cohomology.

Definition 1.7. Let \mathcal{X} be a derived Artin stack and $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ a quasi-coherent complex. For an integer $n \in \mathbf{Z}$, we say that \mathcal{F} is *n -connective* if $\pi_i(u^*\mathcal{F}) = 0$ for all $i < n$ and smooth morphisms $u : X \rightarrow \mathcal{X}$ with X affine. We say \mathcal{F} is *n -coconnective* if $\pi_i(u^*\mathcal{F}) = 0$ for all $i > n$ and smooth morphisms

$u : X \rightarrow \mathcal{X}$ with X affine. The (0-)connective and (0-)coconnective complexes define full subcategories

$$\mathbf{D}_{\mathrm{qc}}(\mathcal{X})_{\geq 0} \subseteq \mathbf{D}_{\mathrm{qc}}(\mathcal{X}), \quad \mathbf{D}_{\mathrm{qc}}(\mathcal{X})_{\leq 0} \subseteq \mathbf{D}_{\mathrm{qc}}(\mathcal{X}),$$

respectively, which together form a canonical t-structure on $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$. This t-structure restricts to $\mathbf{D}_{\mathrm{coh}}(\mathcal{X})$, but typically not to $\mathbf{D}_{\mathrm{perf}}(\mathcal{X})$ unless \mathcal{X} is a classical regular stack. We write

$$\mathrm{Qcoh}(\mathcal{X}) := \mathbf{D}_{\mathrm{qc}}(\mathcal{X})^{\heartsuit}, \quad \mathrm{Coh}(\mathcal{X}) := \mathbf{D}_{\mathrm{coh}}(\mathcal{X})^{\heartsuit}$$

for the hearts. Note that these are insensitive to the derived structure on \mathcal{X} : they are equivalent to the abelian categories of quasi-coherent and coherent sheaves, respectively, on the classical truncation $\mathcal{X}_{\mathrm{cl}}$.

Remark 1.8. Let \mathcal{X} be a derived Artin stack. Suppose that \mathcal{X} has *bounded structure sheaf*, i.e., that the quasi-coherent complex $\mathcal{O}_{\mathcal{X}}$ is n -coconnective for some $n \gg 0$. In that case there is an inclusion $\mathbf{D}_{\mathrm{perf}}(\mathcal{X}) \subseteq \mathbf{D}_{\mathrm{coh}}(\mathcal{X})$. Indeed the properties of perfectness and coherence are both local by definition so this follows from Lemma 1.4.

Remark 1.9. The t-structure on $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$ is left- and right-complete. That is, for every quasi-coherent complex $\mathcal{F} \in \mathbf{D}_{\mathrm{qc}}(\mathcal{X})$, the canonical morphisms

$$\begin{aligned} \mathcal{F} &\rightarrow \varprojlim_{n \geq 0} \tau_{\leq n}(\mathcal{F}), \\ \varinjlim_{n \geq 0} \tau_{\geq -n}(\mathcal{F}) &\rightarrow \mathcal{F} \end{aligned}$$

are invertible. Here $\tau_{\leq n} : \mathbf{D}_{\mathrm{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\mathrm{qc}}(\mathcal{X})_{\leq n}$ and $\tau_{\geq -n} : \mathbf{D}_{\mathrm{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\mathrm{qc}}(\mathcal{X})_{\geq -n}$ are the truncation functors, left and right adjoint to the respective inclusions. This follows from the fact that $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$ is a limit of stable ∞ -categories with left- and right-complete t-structures, and t-exact transition functors (see [Lu3, Prop. 7.1.1.13] and [GR, Chap. 3, Lem. 1.5.8]).

1.3. Tensor, Hom, and inverse/direct images. We have the following basic operations on $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$.

1.3.1. For every derived Artin stack \mathcal{X} , there is a (derived) tensor product $\otimes^{\mathbf{L}}$ on $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$. It is left adjoint, as a bifunctor, to the internal Hom functor $\mathbf{R}\underline{\mathrm{Hom}}$. These are part of a closed symmetric monoidal structure on $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$.

For every fixed perfect complex $\mathcal{F} \in \mathbf{D}_{\mathrm{perf}}(\mathcal{X})$, the operation

$$\mathcal{G} \mapsto \mathcal{F} \otimes^{\mathbf{L}} \mathcal{G}$$

preserves perfectness and coherence in $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$. (One easily reduces to the case where $\mathcal{X} = X$ is affine and $\mathcal{F} = \mathcal{O}_X$.)

Remark 1.10. The symmetric monoidal structure allows the following characterization of perfect complexes over arbitrary \mathcal{X} : namely, $\mathbf{D}_{\mathrm{perf}}(\mathcal{X})$ consists of precisely those objects of $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$ (or of $\mathbf{D}_{\mathrm{coh}}(\mathcal{X})$) which are *dualizable* with respect to the symmetric monoidal structure. This follows by general nonsense (see [Lu3, Prop. 4.6.1.11]) from the case of affines, which is a straightforward exercise (e.g. [Lu4, Prop. 6.2.6.2]).

1.3.2. Given a morphism of derived Artin stacks $f : \mathcal{X} \rightarrow \mathcal{Y}$, there is an inverse image functor

$$\mathbf{L}f^* : \mathbf{D}_{\text{qc}}(\mathcal{Y}) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{X})$$

which is symmetric monoidal. It is right t-exact, i.e., preserves connectivity. It also preserves perfect complexes and pseudocoherent complexes. If f is flat, then $\mathbf{L}f^* = f^*$ is also left t-exact.

1.3.3. The inverse image functor admits a right adjoint, the direct image functor

$$\mathbf{R}f_* : \mathbf{D}_{\text{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{Y}).$$

By adjunction, it is left t-exact, i.e., preserves coconnectivity. If it is affine, then $\mathbf{R}f_* = f_*$ is also right t-exact. If f is representable and proper (i.e., if it is representable, separated of finite type, and satisfies the valuative criterion), then $\mathbf{R}f_*$ preserves pseudocoherent and coherent complexes. See [Lu4, Thms. 5.6.0.2], and Subsect. 1.6 for a discussion of the non-representable case.

Remark 1.11. When f is representable by qcqs derived algebraic spaces, $\mathbf{R}f_*$ commutes with colimits and satisfies base change and projection formulas. That is:

- (i) The functor $\mathbf{L}f^*$ is compact, i.e., its right adjoint $\mathbf{R}f_*$ preserves colimits. See [Lu4, Cor. 3.4.2.2 (2)].
- (ii) For every homotopy cartesian square of derived Artin stacks

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{g} & \mathcal{Y}' \\ \downarrow p & & \downarrow q \\ \mathcal{X} & \xrightarrow{f} & \mathcal{Y}, \end{array}$$

there is a canonical isomorphism

$$\mathbf{L}q^* \mathbf{R}f_* \rightarrow \mathbf{R}g_* \mathbf{L}p^*$$

of functors $\mathbf{D}_{\text{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{Y}')$. See [Lu4, Cor. 3.4.2.2 (3)].

- (iii) For every $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ and $\mathcal{G} \in \mathbf{D}_{\text{qc}}(\mathcal{Y})$, there is a canonical isomorphism

$$\mathbf{R}f_*(\mathcal{F}) \overset{\mathbf{L}}{\otimes} \mathcal{G} \rightarrow \mathbf{R}f_*(\mathcal{F} \overset{\mathbf{L}}{\otimes} \mathbf{L}f^*(\mathcal{G}))$$

which is natural in \mathcal{F} and \mathcal{G} . In other words, $\mathbf{R}f_*$ is $\mathbf{D}_{\text{qc}}(\mathcal{Y})$ -linear when $\mathbf{D}_{\text{qc}}(\mathcal{X})$ is regarded as a $\mathbf{D}_{\text{qc}}(\mathcal{Y})$ -module category via the symmetric monoidal functor $\mathbf{L}f^*$. See [Lu4, Rem. 3.4.2.6].

As we will see in the next subsection, representability can be considerably weakened.

1.4. Finiteness of Tor-amplitude. Recall that a morphism $A \rightarrow B$ of derived commutative rings is of Tor-amplitude $\leq n$ if B is of Tor-amplitude $\leq n$ as a complex over A (Definition 1.3). This condition is fpqc-local on the source and target, so we may extend to derived Artin stacks in the usual manner. For example, a morphism is of Tor-amplitude ≤ 0 if and only if it is flat. We also say a morphism is of *finite Tor-amplitude* if it is of Tor-amplitude $\leq n$ for some $n \geq 0$. We clearly have:

Proposition 1.12. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of derived Artin stacks. If f is of Tor-amplitude $\leq n$, then $\mathbf{L}f^*$ restricts to a functor*

$$\mathbf{L}f^* : \mathbf{D}_{\text{qc}}(\mathcal{Y})_{\leq 0} \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{X})_{\leq n},$$

i.e., it sends coconnective complexes to n -coconnective complexes.

Remark 1.13. Recall that $\mathbf{L}f^*$ always preserves pseudocoherent complexes. Thus when f is of finite Tor-amplitude, it also preserves coherent complexes.

Remark 1.14. If f is proper representable, locally almost of finite presentation, and of finite Tor-amplitude, then $\mathbf{R}f_*$ preserves perfect complexes. See [Lu4, Thm. 6.1.3.2].

Recall that a morphism of derived Artin stacks $f : \mathcal{X} \rightarrow \mathcal{Y}$ is *quasi-smooth* if it is locally of finite presentation and the relative cotangent complex $\mathcal{L}_{\mathcal{X}/\mathcal{Y}}$ is of Tor-amplitude ≤ 1 . Equivalently, it factors smooth-locally on the source as the inclusion $\mathcal{X} \hookrightarrow \mathcal{Y}'$ of the derived zero locus of a section of a vector bundle over a stack \mathcal{Y}' which is smooth over \mathcal{Y} ; see [KhRy, Prop. 2.3.14]. This is a derived version of the notion of *local complete intersection* morphism.

Lemma 1.15. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-smooth morphism of derived Artin stacks. Then f is of finite Tor-amplitude.*

Proof. If f is smooth, then it is of Tor-amplitude ≤ 0 . Therefore we may assume that f is the inclusion of the derived zero locus $\mathcal{X} = X$ of n functions g_1, \dots, g_n on an affine derived scheme $\mathcal{Y} = Y$. We claim that f is of Tor-amplitude $\leq n$. By induction, we may assume that $n = 1$. Then the exact triangle

$$\mathcal{O}_Y \xrightarrow{g_1} \mathcal{O}_Y \rightarrow f_*(\mathcal{O}_X)$$

shows that f is of Tor-amplitude ≤ 1 . □

1.5. Finiteness of cohomological dimension. The functor f_* is well-behaved for many non-representable morphisms as well. The relevant condition is as follows.

Definition 1.16. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of derived Artin stacks. We say that f is of *cohomological dimension $\leq n$* if $\mathbf{R}f_*$ restricts to a functor

$$\mathbf{R}f_* : \mathbf{D}_{\text{qc}}(\mathcal{X})_{\geq 0} \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{Y})_{\geq -n},$$

i.e., it sends connective complexes to $(-n)$ -connective complexes. Equivalently, $\mathbf{R}f_(\mathcal{F})$ is $(-n)$ -connective for every discrete $\mathcal{F} \in \text{Qcoh}(\mathcal{X})$ (see [HLP, Lem. A.1.6]). We say that f is of *finite cohomological dimension* if it is of cohomological dimension $\leq n$ for some integer $n \in \mathbf{N}$.*

Example 1.17. We say that a derived Artin stack \mathcal{X} is of cohomological dimension $\leq n$ if the structural morphism $f : \mathcal{X} \rightarrow \mathrm{Spec}(\mathbf{Z})$ is. Note that this is the case if and only if the complex

$$\mathbf{R}\Gamma(\mathrm{Spec}(\mathbf{Z}), \mathbf{R}f_*(\mathcal{F})) \simeq \mathbf{R}\Gamma(\mathcal{X}, \mathcal{F})$$

is $(-n)$ -connective for all quasi-coherent sheaves $\mathcal{F} \in \mathrm{Qcoh}(\mathcal{X})$. In other words, if

$$H^i(\mathcal{X}, \mathcal{F}) = \pi_{-i}(\mathbf{R}\Gamma(\mathcal{X}, \mathcal{F})) = 0$$

for all $i > n$ and all $\mathcal{F} \in \mathrm{Qcoh}(\mathcal{X})$.

Definition 1.18. We say that a morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ is *universally of finite cohomological dimension* if for every morphism $\mathcal{Y}' \rightarrow \mathcal{Y}$ with \mathcal{Y}' qcqs, the base change $f' : \mathcal{X} \times_{\mathcal{Y}} \mathcal{Y}' \rightarrow \mathcal{Y}'$ is of finite cohomological dimension.

In [HR2], qcqs morphisms that are universally of finite cohomological dimension are called “concentrated”.

Remark 1.19. If \mathcal{Y} is quasi-compact with quasi-affine diagonal, then any qcqs morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ of finite cohomological dimension is also *universally* of finite cohomological dimension. See [HR2, Lem. 2.5(v)].

Our main interest in this condition is the good behaviour of $\mathbf{R}f_*$ it implies (compare Remark 1.11):

Theorem 1.20. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-compact quasi-separated morphism of derived Artin stacks. If f is universally of finite cohomological dimension, then $\mathbf{R}f_*$ commutes with colimits and satisfies the base change and projection formulas (as in Remark 1.11).*

Proof. See [HR2, Thm. 2.6] and [HLP, Prop. A.1.5]. □

Remark 1.21. The fact that $\mathbf{R}f_*$ commutes with colimits (i.e., that $\mathbf{L}f^*$ is a compact functor) implies by adjunction that $\mathbf{L}f^*$ preserves compact objects. In particular, Theorem 1.20 implies that if \mathcal{X} is a qcqs derived Artin stack which is universally of finite cohomological dimension, then the structure sheaf $\mathcal{O}_{\mathcal{X}}$ is compact as an object of $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$.

Remark 1.22. A morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ is of finite cohomological dimension if and only if the underlying morphism $f_{\mathrm{cl}} : \mathcal{X}_{\mathrm{cl}} \rightarrow \mathcal{Y}_{\mathrm{cl}}$ of classical truncations is of finite cohomological dimension. Indeed, this follows by consideration of the following commutative square

$$\begin{array}{ccc} \mathrm{Qcoh}(\mathcal{X}_{\mathrm{cl}}) & \xrightarrow{\mathbf{R}f_{\mathrm{cl},*}} & \mathbf{D}(\mathcal{Y}_{\mathrm{cl}})_{\leq 0} \\ \downarrow i_{\mathcal{X},*} & & \downarrow i_{\mathcal{Y},*} \\ \mathrm{Qcoh}(\mathcal{X}) & \xrightarrow{\mathbf{R}f_*} & \mathbf{D}(\mathcal{Y})_{\leq 0}, \end{array}$$

where the vertical arrows are direct image along the inclusions of the classical truncations. Recall that $i_{\mathcal{X},*}$ and $i_{\mathcal{Y},*}$ are t-exact (since $i_{\mathcal{X}}$ and $i_{\mathcal{Y}}$ are affine morphisms) and that the left-hand arrow is an equivalence (Definition 1.7).

Example 1.23. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of derived Artin stacks. If f is representable by qcqs derived algebraic spaces, then it is universally of finite cohomological dimension. See [Lu4, Prop. 2.5.4.4, Thm. 3.4.2.1].

Example 1.24. Let G be an affine group scheme of finite presentation over an affine scheme S and consider the morphism $f : BG \rightarrow S$, the structural morphism of the classifying stack.

- (i) If G is linearly reductive, then f is universally of cohomological dimension zero.
- (ii) If S is the spectrum of a field of characteristic zero, then f is universally of finite cohomological dimension (see [HR1, Thm. 1.4]).

Example 1.25. Let \mathcal{X} be a qcqs derived 1-Artin stack. Then \mathcal{X} is of finite cohomological dimension under any one of the following conditions:

- (i) \mathcal{X} is of characteristic zero and its stabilizers at all points are affine.
- (ii) \mathcal{X} is of positive characteristic and its stabilizers at all points are linearly reductive.
- (iii) \mathcal{X} is of mixed characteristic and its stabilizers at all points are “nice” (extensions of tame finite étale groups by groups of multiplicative type).
- (iv) \mathcal{X} is of mixed characteristic, its (classical) inertia stack is of finite presentation over \mathcal{X}_{cl} , its stabilizers at all points are affine, and its stabilizers at points of positive characteristic are linearly reductive.

Similarly, a morphism of qcqs derived Artin stacks $f : \mathcal{X} \rightarrow \mathcal{Y}$ is of finite cohomological dimension if its fibres $\mathcal{X} \times_{\mathcal{Y}} Y$ satisfy one of the above conditions, for all morphisms $v : Y \rightarrow \mathcal{Y}$ with Y affine. See [DG, Thm. 1.4.2] and [HR1, Thm. 2.1].

1.6. Cohomological properness. We would now like to study the question of when the direct image functor preserves coherent complexes.

Definition 1.26. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of derived Artin stacks. Assume that f is of finite cohomological dimension. We say that f is *cohomologically proper* if for every coherent sheaf $\mathcal{F} \in \text{Coh}(\mathcal{X})$, the direct image $\mathbf{R}f_*(\mathcal{F}) \in \mathbf{D}_{\text{qc}}(\mathcal{Y})$ is a coherent complex.

Remark 1.27. Since f is of finite cohomological dimension, $\mathbf{R}f_*(\mathcal{F}) \in \mathbf{D}_{\text{qc}}(\mathcal{Y})$ is bounded (cohomologically) for any coherent sheaf $\mathcal{F} \in \text{Coh}(\mathcal{X})$. Thus the condition of Definition 1.26 is that it has coherent cohomologies, i.e., that for every smooth morphism $u : Y \rightarrow \mathcal{Y}$ with Y affine, the groups

$$\mathrm{H}^i(Y, u^* \mathbf{R}f_*(\mathcal{F})) = \pi_{-i}(\mathbf{R}\Gamma(Y, u^* \mathbf{R}f_*(\mathcal{F})))$$

are finitely generated over $\mathrm{H}^0(Y, \mathcal{O}_Y)$ for all i .

Remark 1.28. If $f : \mathcal{X} \rightarrow \mathcal{Y}$ is cohomologically proper, then any coherent complex $\mathcal{F} \in \mathbf{D}_{\text{coh}}(\mathcal{X})$ also has coherent direct image $\mathbf{R}f_*(\mathcal{F}) \in \mathbf{D}_{\text{coh}}(\mathcal{Y})$.

This follows by induction on the Postnikov tower, i.e., using the exact triangles

$$\pi_n(\mathcal{F})[n] \rightarrow \tau_{\leq n}(\mathcal{F}) \rightarrow \tau_{\leq n-1}(\mathcal{F})$$

coming from the (bounded) t-structure on $\mathbf{D}_{\text{coh}}(\mathcal{X})$.

Remark 1.29. As in Remark 1.22, the condition of cohomological properness may be checked on classical truncations.

Proposition 1.30. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of finite cohomological dimension between noetherian derived 1-Artin stacks. If f is proper (of finite type, separated and satisfying the valuative criterion), then it is cohomologically proper.*

Proof. See [Fa, Thm. 1], [Ol, Thm. 1.2]. □

Proposition 1.31. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a cohomologically proper morphism of derived Artin stacks that is universally of finite cohomological dimension. If f is moreover almost of finite presentation and of finite Tor-amplitude, then $\mathbf{R}f_*$ also preserves perfect complexes.*

Proof. Since f is universally of finite cohomological dimension, formation of $\mathbf{R}f_*$ is stable under base change (Theorem 1.20). Since perfectness is smooth-local by definition, we may assume that $\mathcal{Y} = Y$ is affine. By Lemma 1.4(iii) it will suffice to show that the coherent complex $\mathbf{R}f_*(\mathcal{F}) \in \mathbf{D}_{\text{coh}}(Y)$ is of finite Tor-amplitude for every perfect complex $\mathcal{F} \in \mathbf{D}_{\text{perf}}(\mathcal{X})$. Now the claim follows from [Lu4, Prop. 6.1.3.1]. □

1.7. The resolution property. We make a short digression to record a very useful property of stacks, studied in detail in [T2, Tot1, Gr]. For our purposes it will also be useful to consider a stronger “derived” variant of the definition.

Definition 1.32. Let \mathcal{X} be a qcqs derived Artin stack.

- (i) We say that \mathcal{X} has the *resolution property* if, for every quasi-coherent sheaf $\mathcal{F} \in \text{Qcoh}(\mathcal{X})$, there exists a small collection $\{\mathcal{E}_\alpha\}_\alpha$ of finite locally free sheaves on \mathcal{X} and a morphism

$$\phi : \bigoplus_{\alpha} \mathcal{E}_\alpha \rightarrow \mathcal{F}$$

which is surjective (on π_0).

- (ii) We say that \mathcal{X} has the *derived resolution property* if the above condition holds for every quasi-coherent complex $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$.

Remark 1.33. It suffices to check the derived resolution property for connective complexes $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})_{\geq 0}$ only, since the connective cover $\tau_{\geq 0}(\mathcal{F}) \rightarrow \mathcal{F}$ is bijective on π_0 for any $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$.

Remark 1.34. If \mathcal{X} admits the (resp. derived) resolution property and is quasi-compact, and $\mathcal{F} \in \text{Qcoh}(\mathcal{X})$ is of finite type (resp. $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ has $\pi_0(\mathcal{F})$ of finite type), then there exists a surjection $\mathcal{E} \twoheadrightarrow \mathcal{F}$ where \mathcal{E} is a single finite locally free sheaf.

Proposition 1.35. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ is a quasi-projective morphism of derived Artin stacks. If \mathcal{Y} has the (resp. derived) resolution property, then so does \mathcal{X} .*

Proof. See [KhRa, Lem. A.8] for the usual resolution property; the argument for the derived version is the same. \square

Example 1.36. Let G be a *linearly reductive* affine group scheme over an affine scheme S . If BG has the resolution property (e.g., if G is embeddable [AHR2, Rem. 2.5]), then it also has the derived resolution property. Indeed, let $\mathcal{F} \in \mathbf{D}_{\text{qc}}(BG)_{\geq 0}$ be a connective quasi-coherent complex. By the resolution property there exists a surjection $\phi_0 : \bigoplus_{\alpha} \mathcal{E}_{\alpha} \twoheadrightarrow \pi_0(\mathcal{F})$ for some small collection of finite locally free sheaves \mathcal{E}_{α} on BG . Note that each \mathcal{E}_{α} is projective as an object of $\mathbf{D}_{\text{qc}}(BG)_{\geq 0}$, since the underlying object of $\mathbf{D}_{\text{qc}}(S)_{\geq 0}$ is projective and BG is cohomologically affine (see e.g. [Ho2, Lem. 2.17]). Therefore, there exists a lift $\phi : \bigoplus_{\alpha} \mathcal{E}_{\alpha} \twoheadrightarrow \mathcal{F}$ of ϕ_0 , still surjective on π_0 .

Example 1.37. Let S be the spectrum of a field of characteristic zero and let G be an affine group scheme of finite type over S . Then BG has the derived resolution property. Indeed, G is a closed subgroup of $\text{GL}_{n,S}$ for some n . Since the latter is linearly reductive (since S is of characteristic zero), Example 1.36 implies that $B\text{GL}_n$ has the derived resolution property. By Proposition 1.35 it will therefore suffice to show that the morphism $BG \rightarrow B\text{GL}_{n,S}$ is quasi-projective. Indeed, $[\text{GL}_{n,S}/G]$ admits a $\text{GL}_{n,S}$ -equivariant immersion $[\text{GL}_{n,S}/G] \hookrightarrow \mathbf{P}_S(\mathcal{E})$ over S for some finite locally free sheaf \mathcal{E} on BG (see the proof of [Spr, Cor. 5.5.6]), so that $BG \rightarrow B\text{GL}_{n,S}$ is the quotient by $\text{GL}_{n,S}$ of the quasi-projective morphism $[\text{GL}_{n,S}/G] \rightarrow S$.

1.8. Perfect stacks.

Definition 1.38. Let \mathcal{X} be a derived Artin stack. We say that \mathcal{X} is *perfect* if the canonical functor

$$\text{Ind}(\mathbf{D}_{\text{perf}}(\mathcal{X})) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{X})$$

is an equivalence.

In other words, \mathcal{X} is perfect if $\mathbf{D}_{\text{qc}}(\mathcal{X})$ is compactly generated, and the compact objects are precisely the perfect complexes. See [GR, Chap. 1, §7] for background on compactly generated stable ∞ -categories and ind-completion.

Remark 1.39. Various definitions of perfectness can be found in the literature. For example, [BFN, Def. 3.2] and [Lu4, Def. 9.4.4.1] require the further condition that \mathcal{X} has affine or quasi-affine diagonal, respectively.

Theorem 1.40. *Let \mathcal{X} be a qcqs derived 1-Artin stack whose stabilizer at every point is nice (i.e., an extension of a finite étale group scheme, of order prime to the residue characteristics of S , by a multiplicative type group scheme). Then \mathcal{X} is perfect.*

This result was proven jointly with C. Ravi, and is stated in [KhRa, Thm. 2.24] under the assumption that \mathcal{X} has separated diagonal. This assumption was not used in the proof, which we will review below in Subject. 1.9.

Example 1.41. Here are some special cases of Theorem 1.40:

- (i) *Derived algebraic spaces.* Every qcqs derived algebraic space is perfect. This case was first proven by Toën [To4] (see also [Lu4, Prop. 9.6.1.1]), building on previous work like [TT] and [BVdB, Thm. 3.1.1] for classical schemes.
- (ii) *Quotient stacks.* Quotients of qcqs derived algebraic spaces by nice group schemes have nice stabilizers (since subgroups of nice groups are nice), hence are perfect. This was proven in [KrRa, Prop. 3.3] (based on [HR2, HR1]).
- (iii) *Deligne–Mumford stacks and tame Artin stacks.* Any tame qcqs derived Deligne–Mumford stack is perfect. (Recall that tameness means that the stabilizers at all points, which are finite by assumption, are of order coprime to the residue characteristics of the base and hence nice.) This was proven in [To4, Thm. 4.7] in the case of finite inertia, and in the classical case it follows by combining [HR2, Thm. A] and [HR1, Thm. C]. In fact, the latter shows more generally that tame Artin stacks in the sense of [AOV] are perfect.
- (iv) *Classical Artin stacks with nice stabilizers.* The special case of Theorem 1.40 for \mathcal{X} classical was proven under additional assumptions in [AHR1, Thm. 5.1], [AHR2, Prop. 14.1], and [HoKr, Cor. 3.17] (the latter based on [AHHR] and [HR2, Thm. C]).

We also have the following result for quotient stacks:

Theorem 1.42. *Let $\mathcal{X} = [X/G]$ be the quotient of a quasi-projective derived scheme X by a linear action of an affine group scheme G of finite presentation over an affine scheme S . If S is the spectrum of a field of characteristic zero, or if G is linearly reductive and embeddable, then \mathcal{X} is perfect.*

These cases were first proven in [BFN, Cor. 3.22] and [KhRa, Thm. 2.24, A.9.1(a)], respectively.

1.9. Perfectness criteria. In this section we give some criteria to check perfectness of stacks, most notably Theorem 1.50. We then use these to prove Theorems 1.40 and 1.42. First note the following:

Lemma 1.43. *Let \mathcal{X} be a derived Artin stack.*

- (i) *If \mathcal{X} is quasi-separated then every compact object of $\mathbf{D}_{\text{qc}}(\mathcal{X})$ is a perfect complex.*
- (ii) *If $\mathcal{O}_{\mathcal{X}} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ is compact, then every perfect complex on \mathcal{X} is a compact object of $\mathbf{D}_{\text{qc}}(\mathcal{X})$.*

Proof. For (i), let $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ be a compact object. It is enough to show that $u^*(\mathcal{F}) \in \mathbf{D}_{\text{qc}}(X)$ is perfect for every smooth morphism $u: X \rightarrow \mathcal{X}$ with X affine. Since \mathcal{X} is quasi-separated, u is qcqs (and representable), so u^* is a compact functor (Theorem 1.20, Example 1.23). In particular, $u^*(\mathcal{F})$ is compact and hence also perfect, since X is affine.

For (ii), note that for any $\mathcal{F} \in \mathbf{D}_{\text{perf}}(\mathcal{X})$ we have

$$\text{Maps}_{\mathbf{D}_{\text{qc}}(\mathcal{X})}(\mathcal{F}, -) \simeq \text{Maps}_{\mathbf{D}_{\text{qc}}(\mathcal{X})}(\mathcal{O}_{\mathcal{X}}, \mathcal{F}^{\vee} \otimes^{\mathbf{L}} -),$$

where $\mathcal{F}^{\vee} = \mathbf{R}\underline{\text{Hom}}(\mathcal{F}, \mathcal{O}_{\mathcal{X}})$ is the derived dual (see Remark 1.10). Since $\otimes^{\mathbf{L}}$ commutes with colimits in each argument and $\mathcal{O}_{\mathcal{X}}$ is compact, it follows that this functor commutes with colimits, i.e., that \mathcal{F} is compact. \square

Lemma 1.44. *Let \mathcal{X} be a qcqs derived Artin stack. Then \mathcal{X} is perfect if and only if the following conditions hold:*

- (i) *The structure sheaf $\mathcal{O}_{\mathcal{X}}$ is a compact object of $\mathbf{D}_{\text{qc}}(\mathcal{X})$.*
- (ii) *For every nonzero quasi-coherent complex $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$, there exists a perfect complex $\mathcal{E} \in \mathbf{D}_{\text{perf}}(\mathcal{X})$ and a nonzero morphism $\phi: \mathcal{E} \rightarrow \mathcal{F}$.*

Proof. By definition, \mathcal{X} is perfect if and only if $\mathbf{D}_{\text{qc}}(\mathcal{X})$ is compactly generated and its compact objects are precisely the perfect complexes. The latter holds precisely when $\mathcal{O}_{\mathcal{X}} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ is compact (one direction is Lemma 1.43(ii), the other is clear since $\mathcal{O}_{\mathcal{X}}$ is always perfect). Compact generation is equivalent by [Lu4, Cor. C.6.3.3] (which applies because $\mathbf{D}_{\text{qc}}(\mathcal{X})$ is presentable) to the assertion that for every nonzero $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$, there exists a compact object $\mathcal{E} \in \mathbf{D}_{\text{perf}}(\mathcal{X})$ and a nonzero morphism $\phi: \mathcal{E} \rightarrow \mathcal{F}$. The claim follows. \square

Corollary 1.45. *Let \mathcal{X} be a qcqs derived Artin stack. If $\mathcal{O}_{\mathcal{X}} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ is compact (e.g. \mathcal{X} is universally of finite cohomological dimension) and \mathcal{X} has the derived resolution property, then \mathcal{X} is perfect.*

Proof. Let us check the condition Lemma 1.44(ii). Let $\mathcal{F} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$ be a nonzero quasi-coherent complex. Replacing \mathcal{F} by a shift if necessary, we may assume that $\pi_0(\mathcal{F}) \neq 0$. Then the claim follows immediately from the derived resolution property (Definition 1.32). \square

Proposition 1.46. *Let $f: \mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-affine (resp. quasi-projective) morphism of qcqs derived Artin stacks. If \mathcal{Y} is perfect (resp. and has the derived resolution property), then \mathcal{X} is perfect.*

Proof. Since f is universally of finite cohomological dimension (Example 1.23), it follows from Remark 1.21 that $\mathbf{L}f^*$ sends the compact object $\mathcal{O}_{\mathcal{Y}} \in \mathbf{D}_{\text{qc}}(\mathcal{Y})$ to the compact object $\mathcal{O}_{\mathcal{X}} \in \mathbf{D}_{\text{qc}}(\mathcal{X})$. It is therefore enough to show that $\mathbf{L}f^*: \mathbf{D}_{\text{qc}}(\mathcal{Y}) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{X})$ generates under colimits. In the quasi-affine case this is clear. In the quasi-projective case we may reduce (when \mathcal{Y} has the derived resolution property) to the case where \mathcal{X} is a projective bundle over \mathcal{Y} , which follows from [Kh4, Thm. 3.3]. Compare [KhRa, A.9.1(a)]. \square

We now move to the formulation of Theorem 1.50. We will require some preliminary definitions.

Definition 1.47. We say that a derived Artin stack \mathcal{X} is *absolutely perfect* if it is perfect and moreover the canonical functor

$$\mathrm{Ind}(\mathbf{D}_{\mathrm{perf}}(\mathcal{X} \text{ on } Z)) \rightarrow \mathbf{D}_{\mathrm{qc}}(\mathcal{X} \text{ on } Z)$$

is an equivalence for every cocompact closed subset $Z \subseteq |\mathcal{X}|$. Here $\mathbf{D}_{\mathrm{qc}}(\mathcal{X} \text{ on } Z)$ is the full subcategory of $\mathbf{D}_{\mathrm{qc}}(\mathcal{X})$ spanned by quasi-coherent complexes \mathcal{F} supported on Z (i.e., $j^*(\mathcal{F}) = 0$ for $j: \mathcal{X} \setminus Z \hookrightarrow \mathcal{X}$ the complementary open immersion), and similarly for $\mathbf{D}_{\mathrm{perf}}(\mathcal{X} \text{ on } Z)$.

In many cases, perfection automatically implies absolute perfection:

Lemma 1.48. *Let \mathcal{X} be a qcqs derived Artin stack. If \mathcal{X} admits the resolution property, then it is perfect if and only if it is absolutely perfect.*

Proof. Let $Z \subseteq |\mathcal{X}|$ be a cocompact closed subset. By [Ry2, Prop. 8.2] there exists an ideal $\mathcal{I} \subseteq \mathcal{O}_{\mathcal{X}_{\mathrm{cl}}} = \pi_0(\mathcal{O}_{\mathcal{X}})$ of finite type whose vanishing locus is Z . Since \mathcal{X} admits the resolution property, there exists a surjection $\mathcal{E} \twoheadrightarrow \mathcal{I}$ where \mathcal{E} is a finite locally free sheaf on \mathcal{X} . Let $\mathcal{K} \in \mathbf{D}_{\mathrm{qc}}(\mathcal{X})$ denote the Koszul complex of the resulting cosection $\sigma: \mathcal{E} \twoheadrightarrow \mathcal{I} \subseteq \mathcal{O}_{\mathcal{X}}$, i.e., the structure sheaf of its derived zero locus. Note that \mathcal{K} is a perfect complex supported on Z by construction, and hence gives rise to a functor

$$\mathcal{K} \otimes^{\mathbf{L}} -: \mathbf{D}_{\mathrm{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\mathrm{qc}}(\mathcal{X} \text{ on } Z)$$

which sends perfect complexes on \mathcal{X} to perfect complexes supported on Z . It will suffice to show that this functor generates under colimits, or equivalently that its right adjoint $\mathbf{R}\mathrm{Hom}(\mathcal{K}, -) \simeq \mathcal{K}^\vee \otimes -$ is conservative. Since \mathcal{X} is quasi-compact, we may choose a smooth surjection $u: X \twoheadrightarrow \mathcal{X}$ with X affine. Then $u^*: \mathbf{D}_{\mathrm{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\mathrm{qc}}(X)$ is conservative, symmetric monoidal, and sends \mathcal{K} to the Koszul complex of $f^*(\sigma): f^*(\mathcal{E}) \rightarrow \mathcal{O}_X$. Thus we may assume that $\mathcal{X} = X$ is affine, in which case it is enough to show that \mathcal{K} generates $\mathbf{D}_{\mathrm{qc}}(X \text{ on } Z)$ under colimits, which is proven e.g. in [To4, Lem. 4.10]. \square

Recall that an *étale neighbourhood* of a cocompact closed subset $Z \subseteq |\mathcal{X}|$ is an étale morphism $f: \mathcal{X}' \rightarrow \mathcal{X}$ which restricts to an isomorphism $f^{-1}(Z)_{\mathrm{red}} \rightarrow Z_{\mathrm{red}}$. The following, taken from [KhRa, §2.3], is a more flexible variant of [Lu4, Def. 2.5.3.1].

Definition 1.49. Let \mathcal{X} be a quasi-compact derived Artin stack. A *scallop decomposition* $(\mathcal{U}_i, \mathcal{V}_i, u_i)_i$ of \mathcal{X} is a finite filtration by quasi-compact open substacks

$$\emptyset = \mathcal{U}_0 \hookrightarrow \mathcal{U}_1 \hookrightarrow \cdots \hookrightarrow \mathcal{U}_n = \mathcal{X},$$

together with étale neighbourhoods $u_i: \mathcal{V}_i \rightarrow \mathcal{U}_i$ of $\mathcal{U}_i \setminus \mathcal{U}_{i-1}$.

We can now state the criterion, which is abstracted out of the proof of [KhRa, Thm. 2.24]:

Theorem 1.50. *Let \mathcal{X} be a qcqs derived Artin stack. Suppose given a scallop decomposition $(\mathcal{U}_i, \mathcal{V}_i, u_i)_i$ of \mathcal{X} . If each \mathcal{V}_i is absolutely perfect (resp. each $\mathcal{O}_{\mathcal{V}_i}$ is compact), then \mathcal{X} is perfect (resp. $\mathcal{O}_{\mathcal{X}}$ is compact).*

Proof. One argues exactly as in the proof of [KhRa, Thm. 2.24] (which itself is adapted from arguments of [TT] and [BVdB]). Namely, one first shows as in [KhRa, Lem. 2.27] that the compact objects of $\mathbf{D}_{\text{qc}}(\mathcal{X})$ are the perfect complexes. Then one proceeds by an induction on the scallop decomposition to glue together a set of perfect complexes generating $\mathbf{D}_{\text{qc}}(\mathcal{X})$, using descent by étale neighbourhoods for $\mathbf{D}_{\text{qc}}(-)$. \square

Remark 1.51. In fact a small modification of the proof shows that \mathcal{X} is even absolutely perfect, but we will not need this.

Finally we apply these results to prove Theorems 1.40 and 1.42.

Proof of Theorem 1.42. By Examples 1.36 and 1.37, BG has the derived resolution property. By Example 1.24 it is also universally of finite cohomological dimension (Remark 1.21). By Proposition 1.35 and Example 1.23, the same then holds for $\mathcal{X} = [X/G]$. Thus \mathcal{X} is perfect by Corollary 1.45. \square

Proof of Theorem 1.40. Suppose that \mathcal{X} has nice stabilizers. By the local structure theorem of [AHHR] (cf. [KhRa, Thm. 2.12(ii)]), \mathcal{X} admits a scallop decomposition $(\mathcal{U}_i, \mathcal{V}_i, u_i)_i$ where each \mathcal{V}_i is a quotient of a quasi-affine derived scheme by an action of a linearly reductive embeddable group scheme over an affine base. By Theorem 1.42, each \mathcal{V}_i is perfect. Since it also admits the resolution property (Proposition 1.35 and Example 1.36), it is moreover absolutely perfect by Lemma 1.48. Now the claim follows from Theorem 1.50. \square

1.10. Continuity. The following is a generalization of Thomason's [TT, Prop. 3.20] from the case of qcqs schemes.

Theorem 1.52. *Let $(\mathcal{X}_\alpha)_\alpha$ be a cofiltered system of derived Artin stacks with affine transition morphisms. Denote by \mathcal{X} the limit and by $p_\alpha : \mathcal{X} \rightarrow \mathcal{X}_\alpha$ the projections. Then we have:*

- (i) *The inverse image functors p_α^* induce a canonical equivalence of stable presentable ∞ -categories*

$$\varinjlim_{\alpha} \mathbf{D}_{\text{qc}}(\mathcal{X}_\alpha) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{X}),$$

where the colimit is taken in the (very large) ∞ -category of presentable ∞ -categories and colimit-preserving functors.

- (ii) *If \mathcal{X} and each \mathcal{X}_α are perfect, then p_α^* also induce an equivalence*

$$\varinjlim_{\alpha} \mathbf{D}_{\text{perf}}(\mathcal{X}_\alpha) \rightarrow \mathbf{D}_{\text{perf}}(\mathcal{X}),$$

where the colimit is taken in the (large) ∞ -category of small ∞ -categories.

Proof. When \mathcal{X} and \mathcal{X}_α are perfect, the second equivalence is obtained by passage to compact objects from the first (see [Lu1, Prop. 5.5.7.8, Cor. 4.4.5.21]).

The first statement is equivalent by [Lu1, Cor. 5.5.3.4, Thm. 5.5.3.18] to the assertion that the functor

$$\mathbf{D}_{\text{qc}}(\mathcal{X}) \rightarrow \varprojlim_{\alpha} \mathbf{D}_{\text{qc}}(\mathcal{X}_\alpha), \quad (1.53)$$

induced by the direct image functors $p_{\alpha,*}$, is an equivalence (where the limit is taken in the ∞ -category of large ∞ -categories).

Without loss of generality we may assume that the indexing category admits a terminal object 0. Let T_0 be an affine derived scheme and $t_0 : T_0 \rightarrow \mathcal{X}_0$ a smooth morphism, and consider the analogous functor

$$\mathbf{D}_{\text{qc}}(T) \rightarrow \varprojlim_{\alpha \geq \alpha_0} \mathbf{D}_{\text{qc}}(T_\alpha)$$

where $T_\alpha = T_0 \times_{\mathcal{X}_{\alpha_0}} \mathcal{X}_\alpha$ and $T = T_0 \times_{\mathcal{X}_{\alpha_0}} \mathcal{X}$. Since the transition maps $\mathcal{X}_\alpha \rightarrow \mathcal{X}_{\alpha_0}$ are affine, T and each T_α is affine. Therefore by [Lu4, Cor. 4.5.1.5] this functor is an equivalence.

Since the transition maps are affine, the base change formula (Remark 1.11) implies that the diagram

$$\begin{array}{ccccccccccc} \mathbf{D}_{\text{qc}}(\mathcal{X}) & \longrightarrow & \cdots & \longrightarrow & \mathbf{D}_{\text{qc}}(\mathcal{X}_\alpha) & \longrightarrow & \mathbf{D}_{\text{qc}}(\mathcal{X}_\beta) & \longrightarrow & \cdots & \longrightarrow & \mathbf{D}_{\text{qc}}(\mathcal{X}_0) \\ \downarrow & & & & \downarrow & & \downarrow & & & & \downarrow \\ \mathbf{D}_{\text{qc}}(T) & \longrightarrow & \cdots & \longrightarrow & \mathbf{D}_{\text{qc}}(T_\alpha) & \longrightarrow & \mathbf{D}_{\text{qc}}(T_\beta) & \longrightarrow & \cdots & \longrightarrow & \mathbf{D}_{\text{qc}}(T_0) \end{array}$$

commutes. Therefore, the functor (1.53) is the limit over the analogous functors for every pair (T_0, t_0) , which are all equivalences by above. \square

Corollary 1.54. *Let \mathcal{X} be a perfect derived Artin stack. Suppose that \mathcal{X} is the limit of a cofiltered system $(\mathcal{X}_\alpha)_\alpha$ of perfect derived Artin stacks with affine transition morphisms. Then the equivalence of Theorem 1.52(ii) restricts to an equivalence*

$$\varprojlim_{\alpha} \mathbf{D}_{\text{perf}}(\mathcal{X}_\alpha)_{\text{locfr}} \rightarrow \mathbf{D}_{\text{perf}}(\mathcal{X})_{\text{locfr}}$$

where $(-)_{\text{locfr}}$ denotes the full subcategory of locally free sheaves of finite rank.

Proof. Since the property is smooth-local, we may reduce as in the proof of Theorem 1.52 to the case where $\mathcal{X} = X$ and $\mathcal{X}_\alpha = X_\alpha$ are affine. Choose an index 0 and let $\mathcal{F}_0 \in \mathbf{D}_{\text{perf}}(X_0)$, $\mathcal{F}_\alpha = \mathcal{F}_0|_{X_\alpha} \in \mathbf{D}_{\text{perf}}(X_\alpha)$ for every α , and $\mathcal{F} = \mathcal{F}_0|_X \in \mathbf{D}_{\text{perf}}(X)$. Since inverse image preserves finite locally free sheaves, it is enough to show that if \mathcal{F} is finite locally free, then so is \mathcal{F}_α for some α . For every α let $U_\alpha \subseteq X_\alpha$ denote the open locus where \mathcal{F}_α is finite locally free (see [Lu4, Prop. 2.9.3.2]). By construction we have $U_\alpha = U_0 \times_{X_0} X_\alpha$ for all α and $U = U_0 \times_{X_0} X$. As an open subscheme of $X_0 = \text{Spec}(A_0)$, U_0 can be written as the union of principal opens $U_{0,i} = \text{Spec}(A_0[f_i^{-1}])$ for some elements $f_i \in \pi_0(A_0)$. By base change, the opens $U_{0,i} \times_{X_0} X = \text{Spec}(A[f_i^{-1}])$

then cover $X = \text{Spec}(A)$, i.e., we can write $1 \in \pi_0(A)$ as a linear combination of the images of f_i . But then the same must hold in $\pi_0(A_\alpha)$ for some large enough α . That is, $U_\alpha = X_\alpha$, so that \mathcal{F}_α is finite locally free. \square

2. ALGEBRAIC K-THEORY

For a commutative ring R , Quillen's algebraic K-theory spectrum $K(R)$ (see [Q2]) can be defined succinctly as the homotopy-theoretic group completion [Q1, Ma, GGN] of the groupoid of finitely generated projective R -modules, regarded as an \mathcal{E}_∞ -monoid under direct sum. This can be viewed either as an \mathcal{E}_∞ -group or a connective spectrum. This description is also valid for a derived commutative ring R (see [Lu2, Lect. 19, Thm. 5]).

For a scheme X , doing the same construction with the groupoid of finite locally free \mathcal{O}_X -modules (or vector bundles over X of finite rank) gives rise to Quillen's original construction of $K(X)$. The Thomason–Trobaugh definition [TT] is more sophisticated: it replaces vector bundles by perfect complexes and requires Waldhausen's S_\bullet -construction instead group completion. The advantage of Thomason–Trobaugh K-theory is its excellent behaviour over arbitrary (qcqs) schemes: for instance, there are localization and Mayer–Vietoris long exact sequences.

Thomason–Trobaugh K-theory and Quillen K-theory agree in the presence of the resolution property (Subsect. 1.7), which almost always holds for schemes (e.g. any quasi-projective scheme admits the resolution property). For stacks however the resolution property essentially amounts to being a global quotient [Tot1, Gr]. The Thomason–Trobaugh construction becomes therefore even more crucial in the world of stacks.

2.1. Definitions. Let \mathcal{A} be a stable ∞ -category. The Waldhausen S_\bullet -construction makes sense in this setting and can be used to produce the K-theory space of \mathcal{A} . Iterating it gives rise to an \mathcal{E}_∞ -group structure on this space, and hence to a connective spectrum $K(\mathcal{A})$. See [Lu3, Rem. 1.2.2.5], [BGT, Sect. 7], and [Ba2, Sect. 10]. Using a generalization of the Bass construction (see [CiKh, Sect. 4]), one can also produce a Bass–Thomason–Trobaugh K-theory spectrum $K^B(\mathcal{A})$ whose connective cover $K^B(\mathcal{A})_{\geq 0}$ is $K(\mathcal{A})$. Its homotopy groups are denoted

$$K_i(\mathcal{A}) = \pi_i(K^B(\mathcal{A})), \quad i \in \mathbf{Z},$$

so that $K_i(\mathcal{A}) \simeq \pi_i(K(\mathcal{A}))$ for $i \geq 0$. Any exact functor of stable ∞ -categories $\mathcal{A} \rightarrow \mathcal{B}$ induces maps of spectra $K(\mathcal{A}) \rightarrow K(\mathcal{B})$ and $K^B(\mathcal{A}) \rightarrow K^B(\mathcal{B})$.

Theorem 2.1. *Let $j^* : \mathcal{A} \rightarrow \mathcal{B}$ be a functor between compactly generated stable ∞ -categories with fully faithful right adjoint j_* . Suppose that j^* is compact (equivalently, preserves compact objects) and its kernel \mathcal{A}_0 is compactly generated. Then we have:*

- (i) Proto-localization. *There is a fibre sequence of connective spectra*

$$K(\mathcal{A}_0^\omega) \rightarrow K(\mathcal{A}^\omega) \rightarrow K(\mathcal{B}^\omega).$$

(ii) Localization. *There is an exact triangle of spectra*

$$K^{\mathbb{B}}(\mathcal{A}_0^\omega) \rightarrow K^{\mathbb{B}}(\mathcal{A}^\omega) \rightarrow K^{\mathbb{B}}(\mathcal{B}^\omega).$$

In particular, there is a long exact sequence

$$\cdots \xrightarrow{\partial} K_i(\mathcal{A}_0^\omega) \rightarrow K_i(\mathcal{A}^\omega) \rightarrow K_i(\mathcal{B}^\omega) \xrightarrow{\partial} K_{i-1}(\mathcal{A}_0^\omega) \rightarrow \cdots.$$

Proof. See [Ba2, Prop. 10.20] for the first statement. The second follows from the first and [CiKh, Thm. 4.5.7]. \square

Corollary 2.2 (Additivity). *For any semi-orthogonal decomposition of \mathcal{A} into stable subcategories, the spectra $K(\mathcal{A})$ and $K^{\mathbb{B}}(\mathcal{A})$ both split into direct sums.*

Proof. As in [Kh4, Lem. 2.8], we may reduce to the case of a semi-orthogonal decomposition of length 2. Applying Theorem 2.1 to the ind-completions, we get a fibre sequence (resp. exact triangle), which is split by the map $j_* : K(\mathcal{B}) \rightarrow K(\mathcal{A})$ (resp. $j_* : K^{\mathbb{B}}(\mathcal{B}) \rightarrow K^{\mathbb{B}}(\mathcal{A})$). \square

Remark 2.3. If the kernel \mathcal{A}_0 is not compactly generated, one can still identify the fibre in Theorem 2.1 as the Efimov K-theory of \mathcal{A}_0 (see [Ho3]).

Theorem 2.4. *Suppose given a commutative square*

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array}$$

of compactly generated stable ∞ -categories and compact colimit-preserving functors. If the square is cartesian (in the ∞ -category of large ∞ -categories) and g^ has fully faithful right adjoint g_* , then the induced square of spectra*

$$\begin{array}{ccc} K^{\mathbb{B}}(\mathcal{A}^\omega) & \xrightarrow{f^*} & K^{\mathbb{B}}(\mathcal{B}^\omega) \\ \downarrow p^* & & \downarrow q^* \\ K^{\mathbb{B}}(\mathcal{A}'^\omega) & \xrightarrow{g^*} & K^{\mathbb{B}}(\mathcal{B}'^\omega) \end{array}$$

is cartesian.

Proof. The cartesianness of the square implies that f_* is also fully faithful. Since the ∞ -category of spectra is stable, it will suffice to show that the horizontal fibres are isomorphic. If the kernels of f^* and g^* are compactly generated, this follows immediately from Theorem 2.1. By Remark 2.3, this holds even otherwise. (See [Ho3, Cor. 13] and, for another proof, [Ta, Thm. 18].) \square

Theorem 2.5 (Continuity). *Let \mathcal{A} be a filtered colimit of stable ∞ -categories \mathcal{A}_α . Then the canonical maps*

$$\begin{aligned} \lim_{\alpha} \mathrm{K}(\mathcal{A}_\alpha) &\rightarrow \mathrm{K}(\mathcal{A}), \\ \lim_{\alpha} \mathrm{K}^{\mathrm{B}}(\mathcal{A}_\alpha) &\rightarrow \mathrm{K}^{\mathrm{B}}(\mathcal{A}) \end{aligned}$$

are invertible.

Proof. For $\mathrm{K}(-)$ this follows by inspection of the Waldhausen S_\bullet -construction (see e.g. [BGT, Prop. 7.10]). For $\mathrm{K}^{\mathrm{B}}(-)$, note that it is enough to show the map is invertible after taking the n -connective cover $\tau_{\geq n}$ for every n . The induced map is, by construction of $\mathrm{K}^{\mathrm{B}}(-)$ and because $\tau_{\geq n}$ commutes with colimits, a retract of the map

$$\lim_{\alpha} \mathrm{K}(\mathcal{A}_\alpha \otimes \mathcal{C}^{\otimes n}) \rightarrow \mathrm{K}(\mathcal{A} \otimes \mathcal{C}^{\otimes n})$$

which is invertible by the $\mathrm{K}(-)$ statement. Here \mathcal{C} is the stable ∞ -category of perfect complexes over an analogue of the multiplicative group \mathbf{G}_m over the sphere spectrum. See [CiKh, Not. 4.2.4] and compare the proof of [CiKh, Thm. 4.5.7(ii)]. \square

Definition 2.6. For a derived Artin stack \mathcal{X} , write

$$\mathrm{K}(\mathcal{X}) = \mathrm{K}(\mathbf{D}_{\mathrm{perf}}(\mathcal{X})), \quad \mathrm{K}^{\mathrm{B}}(\mathcal{X}) = \mathrm{K}^{\mathrm{B}}(\mathbf{D}_{\mathrm{perf}}(\mathcal{X}))$$

for the Thomason–Trobaugh and Bass–Thomason–Trobaugh K -theory spectra.

From now on we'll mostly restrict our attention to K^{B} , since statements about K can be recovered simply by passing to connective covers.

2.2. Operations. The various operations on perfect complexes give rise to operations in K -theory.

The tensor product on $\mathbf{D}_{\mathrm{perf}}(\mathcal{X})$ induces a cup product

$$\cup : \mathrm{K}^{\mathrm{B}}(\mathcal{X}) \otimes \mathrm{K}^{\mathrm{B}}(\mathcal{X}) \rightarrow \mathrm{K}^{\mathrm{B}}(\mathcal{X})$$

which is part of an \mathcal{E}_∞ -ring structure on $\mathrm{K}^{\mathrm{B}}(\mathcal{X})$.

For any morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$, the inverse image functor $f^* : \mathbf{D}_{\mathrm{perf}}(\mathcal{Y}) \rightarrow \mathbf{D}_{\mathrm{perf}}(\mathcal{X})$ gives rise to a map

$$f^* : \mathrm{K}^{\mathrm{B}}(\mathcal{Y}) \rightarrow \mathrm{K}^{\mathrm{B}}(\mathcal{X}).$$

Since f^* is symmetric monoidal, this map is multiplicative (a homomorphism of \mathcal{E}_∞ -ring spectra).

If f is proper, of finite cohomological dimension, almost of finite presentation, and of finite Tor-amplitude, then there is a Gysin map

$$f_* : \mathrm{K}^{\mathrm{B}}(\mathcal{X}) \rightarrow \mathrm{K}^{\mathrm{B}}(\mathcal{Y}).$$

By Theorem 1.20 this map is $\mathrm{K}^{\mathrm{B}}(\mathcal{Y})$ -linear (a homomorphism of $\mathrm{K}^{\mathrm{B}}(\mathcal{Y})$ -module spectra). In other words:

Proposition 2.7 (Projection formula). *If $f : \mathcal{X} \rightarrow \mathcal{Y}$ is proper, of finite cohomological dimension, almost of finite presentation, and of finite Tor-amplitude, then we have canonical identifications*

$$f_*(x) \cup y \simeq f_*(x \cup f^*(y))$$

in $\mathbf{K}^{\mathbf{B}}(\mathcal{Y})$, for all $x \in \mathbf{K}^{\mathbf{B}}(\mathcal{X})$, $y \in \mathbf{K}^{\mathbf{B}}(\mathcal{Y})$.

We also have (by Theorem 1.20):

Proposition 2.8 (Base change formula). *Suppose given a homotopy cartesian square of derived Artin stacks*

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{g} & \mathcal{Y}' \\ \downarrow p & & \downarrow q \\ \mathcal{X} & \xrightarrow{f} & \mathcal{Y}. \end{array}$$

If f is proper, of finite cohomological dimension, almost of finite presentation, and of finite Tor-amplitude, then we have a canonical homotopy

$$q^* f_* \simeq g_* p^*$$

of maps $\mathbf{K}^{\mathbf{B}}(\mathcal{X}) \rightarrow \mathbf{K}^{\mathbf{B}}(\mathcal{Y}')$.

If $Z \subseteq |\mathcal{X}|$ is a cocompact closed subset, then we write

$$\mathbf{K}^{\mathbf{B}}(\mathcal{X} \text{ on } Z) := \mathbf{K}^{\mathbf{B}}(\mathbf{D}_{\text{perf}}(\mathcal{X} \text{ on } Z))$$

for the Bass–Thomason–Trobaugh K-theory spectrum of $\mathbf{D}_{\text{perf}}(\mathcal{X} \text{ on } Z)$, defined as in Definition 1.47.

Theorem 2.9 (Localization). *If \mathcal{X} is absolutely perfect, then for every cocompact closed subset $Z \subseteq |\mathcal{X}|$ there is a canonical exact triangle of spectra*

$$\mathbf{K}^{\mathbf{B}}(\mathcal{X} \text{ on } Z) \rightarrow \mathbf{K}^{\mathbf{B}}(\mathcal{X}) \xrightarrow{j^*} \mathbf{K}^{\mathbf{B}}(\mathcal{X} \setminus Z).$$

Proof. Since \mathcal{X} is perfect, so is $\mathcal{X} \setminus Z$ (since $j^* : \mathbf{D}_{\text{qc}}(\mathcal{X}) \rightarrow \mathbf{D}_{\text{qc}}(\mathcal{X} \setminus Z)$ is compact and generates under colimits). Hence we have $\mathbf{D}_{\text{perf}}(\mathcal{X}) \simeq \mathbf{D}_{\text{qc}}(\mathcal{X})^\omega$ and similarly for $\mathcal{X} \setminus Z$, as well as $\mathbf{D}_{\text{perf}}(\mathcal{X} \text{ on } Z) \simeq \mathbf{D}_{\text{qc}}(\mathcal{X} \text{ on } Z)^\omega$. Thus we may apply Theorem 2.1. \square

2.3. Projective bundles and blow-ups. Let \mathcal{X} be a derived Artin stack. For any locally free sheaf \mathcal{E} on \mathcal{X} of rank $n + 1$, $n \geq 0$, we may consider the associated projective bundle

$$q : \mathbf{P}_{\mathcal{X}}(\mathcal{E}) \rightarrow \mathcal{X}.$$

This is the universal derived stack over \mathcal{X} equipped with a surjection $q^*(\mathcal{E}) \twoheadrightarrow \mathcal{O}(1)$ onto an invertible sheaf. The morphism q is smooth of relative dimension n , proper and schematic (see [Kh4, Prop. 3.1]).

The following is a generalization of [TT, Thm. 7.3], [T3].

Theorem 2.10. *The maps*

$$\mathrm{K}^{\mathrm{B}}(X) \rightarrow \mathrm{K}^{\mathrm{B}}(\mathbf{P}_{\mathcal{X}}(\mathcal{E})), \quad x \mapsto q^*(x) \cup [\mathcal{O}(-k)],$$

induce an isomorphism

$$\mathrm{K}^{\mathrm{B}}(\mathbf{P}_{\mathcal{X}}(\mathcal{E})) \simeq \bigoplus_{k=0}^n \mathrm{K}^{\mathrm{B}}(\mathcal{X}).$$

Proof. The standard semi-orthogonal decomposition of Orlov–Thomason on $\mathbf{D}_{\mathrm{perf}}(\mathbf{P}_{\mathcal{X}}(\mathcal{E}))$ extends to this setting by [Kh4, Thm. B]. Thus the formula follows from Corollary 2.2. See [Kh4, Cor. 3.6]. \square

For any quasi-smooth closed immersion of derived Artin stacks $i: \mathcal{Z} \rightarrow \mathcal{X}$, say of virtual codimension n , we may form the derived blow-up $\mathrm{Bl}_{\mathcal{Z}} \mathcal{X}$ as in [KhRy]. This fits in a commutative square

$$\begin{array}{ccc} \mathbf{P}_{\mathcal{Z}}(\mathcal{N}_{\mathcal{Z}/\mathcal{X}}) & \xrightarrow{i_D} & \mathrm{Bl}_{\mathcal{Z}} \mathcal{X} \\ \downarrow q & & \downarrow p \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

which is cartesian on classical truncations (but not homotopy cartesian unless $n = 1$). The upper closed immersion i_D is a virtual Cartier divisor, i.e., a quasi-smooth closed immersion of virtual codimension 1. The morphism $q: \mathbf{P}_{\mathcal{Z}}(\mathcal{N}_{\mathcal{Z}/\mathcal{X}}) \rightarrow \mathcal{Z}$ is the projection of the projective bundle associated to the conormal sheaf $\mathcal{N}_{\mathcal{Z}/\mathcal{X}}$. The projection $p: \mathrm{Bl}_{\mathcal{Z}} \mathcal{X} \rightarrow \mathcal{X}$ is quasi-smooth of relative virtual dimension 0, proper, and schematic. The following is a generalization of [T4].

Theorem 2.11. *The maps $p^*: \mathrm{K}^{\mathrm{B}}(\mathcal{X}) \rightarrow \mathrm{K}^{\mathrm{B}}(\mathrm{Bl}_{\mathcal{Z}} \mathcal{X})$ and*

$$\mathrm{K}^{\mathrm{B}}(\mathcal{Z}) \rightarrow \mathrm{K}^{\mathrm{B}}(\mathrm{Bl}_{\mathcal{Z}} \mathcal{X}), \quad x \mapsto i_{D,*}(q^*(x) \cup [\mathcal{O}(-k)]),$$

induce an isomorphism

$$\mathrm{K}^{\mathrm{B}}(\mathrm{Bl}_{\mathcal{Z}} \mathcal{X}) \simeq \mathrm{K}^{\mathrm{B}}(\mathcal{X}) \oplus \bigoplus_{k=1}^{n-1} \mathrm{K}^{\mathrm{B}}(\mathcal{Z}).$$

Proof. The standard semi-orthogonal decomposition of Orlov–Thomason on $\mathbf{D}_{\mathrm{perf}}(\mathrm{Bl}_{\mathcal{Z}} \mathcal{X})$ extends to this setting by [Kh4, Thm. C]. Thus the formula follows from Corollary 2.2. See [Kh4, Cor. 4.4]. \square

Corollary 2.12. *The induced square*

$$\begin{array}{ccc} \mathrm{K}^{\mathrm{B}}(\mathcal{X}) & \xrightarrow{i^*} & \mathrm{K}^{\mathrm{B}}(\mathcal{Z}) \\ \downarrow p^* & & \downarrow q^* \\ \mathrm{K}^{\mathrm{B}}(\mathrm{Bl}_{\mathcal{Z}} \mathcal{X}) & \xrightarrow{i_D^*} & \mathrm{K}^{\mathrm{B}}(\mathbf{P}_{\mathcal{Z}}(\mathcal{N}_{\mathcal{Z}/\mathcal{X}})) \end{array}$$

is cartesian.

Proof. Combine Theorems 2.10 and 2.11. \square

2.4. Excising closed subsets.

Theorem 2.13. *Let \mathcal{X} and \mathcal{X}' be perfect derived Artin stacks. Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a morphism and $Z \subseteq |\mathcal{X}|$ a cocompact closed subset. Suppose one of the following conditions holds:*

- (i) *Formal neighbourhood. The morphism f is representable and is an isomorphism infinitely near Z . That is, the induced morphism $\mathcal{X}'_{f^{-1}(Z)} \rightarrow \mathcal{X}_Z$ on formal completions is invertible.*
- (ii) *Étale neighbourhood. The morphism f is étale and restricts to an isomorphism $f^{-1}(Z)_{\text{red}} \rightarrow Z_{\text{red}}$.*

Then the induced square

$$\begin{array}{ccc} \mathrm{K}^{\mathrm{B}}(\mathcal{X}) & \longrightarrow & \mathrm{K}^{\mathrm{B}}(\mathcal{X} \setminus Z) \\ \downarrow f^* & & \downarrow \\ \mathrm{K}^{\mathrm{B}}(\mathcal{X}') & \longrightarrow & \mathrm{K}^{\mathrm{B}}(\mathcal{X}' \setminus f^{-1}(Z)) \end{array}$$

is cartesian.

Proof. Follows from Theorem 2.4 and the corresponding statement at the level of \mathbf{D}_{qc} , see e.g. [BKRS, Thm. 4.1.1, Rem. 4.1.4]. \square

Corollary 2.14. *Let \mathcal{X} be a perfect derived Artin stack. If $\mathcal{U} \subseteq \mathcal{X}$ and $\mathcal{V} \subseteq \mathcal{X}$ are open substacks such that $\mathcal{X} = \mathcal{U} \cup \mathcal{V}$, then the square*

$$\begin{array}{ccc} \mathrm{K}^{\mathrm{B}}(\mathcal{X}) & \longrightarrow & \mathrm{K}^{\mathrm{B}}(\mathcal{U}) \\ \downarrow & & \downarrow \\ \mathrm{K}^{\mathrm{B}}(\mathcal{V}) & \longrightarrow & \mathrm{K}^{\mathrm{B}}(\mathcal{U} \cap \mathcal{V}) \end{array}$$

is cartesian. In particular, there is a long exact Mayer–Vietoris sequence

$$\cdots \xrightarrow{\partial} \mathrm{K}_i(\mathcal{X}) \rightarrow \mathrm{K}_i(\mathcal{U}) \oplus \mathrm{K}_i(\mathcal{V}) \rightarrow \mathrm{K}_i(\mathcal{U} \cap \mathcal{V}) \xrightarrow{\partial} \mathrm{K}_{i-1}(\mathcal{X}) \rightarrow \cdots.$$

Proof. Apply Theorem 2.13 with $\mathcal{Z} \rightarrow \mathcal{X}$ a closed immersion complementary to \mathcal{U} and $\mathcal{X}' = \mathcal{V}$. \square

Remark 2.15. By [Lu1, Thm. 7.3.5.2], Corollary 2.14 implies that K^{B} satisfies Zariski descent on the site of perfect derived Artin stacks. Similarly Theorem 2.13(ii) implies that it satisfies descent for Nisnevich’s completely decomposed étale topology. Indeed this topology is generated by families $\{f : \mathcal{X}' \rightarrow \mathcal{X}, j : \mathcal{X} \setminus Z \hookrightarrow \mathcal{X}\}$, where f is an étale neighbourhood of a cocompact closed $Z \subseteq |\mathcal{X}|$ and j is the inclusion of the complement (see [HoKr, Prop. 2.9] for the case of stacks, [MV, §3, Prop. 1.4] for the case of noetherian finite-dimensional schemes). Hence the claim follows from [Kh1, Thm. 2.2.7].

2.5. Excising open subsets. The problem of excising open subsets is more subtle. We are given a morphism of derived Artin stacks $f : \mathcal{X}' \rightarrow \mathcal{X}$ and quasi-compact open subsets $\mathcal{U} \subseteq \mathcal{X}$, $\mathcal{U}' \subseteq \mathcal{X}'$. Let $\mathcal{Z} \rightarrow \mathcal{X}$ and $\mathcal{Z}' \rightarrow \mathcal{X}'$ be closed immersions complementary to \mathcal{U} and \mathcal{U}' , respectively, fitting in a commutative square

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{Z} & \longrightarrow & \mathcal{X} \end{array}$$

which is cartesian on underlying classical truncations.

For example, if f is affine and the square is also cocartesian, then it is called a *Milnor square*. Algebraic K-theory does not generally satisfy excision for Milnor squares (see [Sw]), but it does for Milnor squares that are *homotopy cartesian*:

Theorem 2.16. *Suppose given a Milnor square as above consisting of perfect derived Artin stacks. If the square is moreover homotopy cartesian, then the induced square of spectra*

$$\begin{array}{ccc} \mathrm{K}^{\mathrm{B}}(\mathcal{X}) & \longrightarrow & \mathrm{K}^{\mathrm{B}}(\mathcal{Z}) \\ \downarrow & & \downarrow \\ \mathrm{K}^{\mathrm{B}}(\mathcal{X}') & \longrightarrow & \mathrm{K}^{\mathrm{B}}(\mathcal{Z}') \end{array}$$

is cartesian.

Proof. See [LT, Thm. A] in the affine case and [BKRS] for the case of stacks. \square

For a general Milnor square, we can still get an excision statement if we replace the closed substacks by their formal completions. In fact, we have the following refinement where we consider the natural pro-spectra $\widehat{\mathrm{K}}^{\mathrm{B}}(\mathcal{X}_{\mathcal{Z}}^{\wedge})$, $\widehat{\mathrm{K}}^{\mathrm{B}}(\mathcal{X}'_{\mathcal{Z}'}^{\wedge})$ associated to the formal completions (viewed as ind-stacks).

Theorem 2.17. *For any Milnor square as above consisting of noetherian derived 1-Artin stacks with bounded structure sheaves, affine diagonals, and nice stabilizers, the induced square of pro-spectra*

$$\begin{array}{ccc} \{\mathrm{K}^{\mathrm{B}}(\mathcal{X})\} & \longrightarrow & \widehat{\mathrm{K}}^{\mathrm{B}}(\mathcal{X}_{\mathcal{Z}}^{\wedge}) \\ \downarrow & & \downarrow \\ \{\mathrm{K}^{\mathrm{B}}(\mathcal{X}')\} & \longrightarrow & \widehat{\mathrm{K}}^{\mathrm{B}}(\mathcal{X}'_{\mathcal{Z}'}^{\wedge}) \end{array}$$

is cartesian.

Proof. See [BKRS]. In the affine case see also [LT, Thm. 2.32] where it is proven that the square is “weakly cartesian”. \square

Another interesting case is that of a proper representable morphism $f : \mathcal{X}' \rightarrow \mathcal{X}$ which restricts to an isomorphism $\mathcal{U}' \rightarrow \mathcal{U}$. Then the square is often called a *proper cdh square* or *abstract blow-up square*. K-theory typically doesn't satisfy excision with respect to such squares either, but again it holds if we pass to formal completions.

Theorem 2.18 (Proper excision). *Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a proper representable morphism of noetherian derived 1-Artin stacks with bounded structure sheaves, affine diagonal, and nice stabilizers. Let $\mathcal{U} \subseteq \mathcal{X}$ and $\mathcal{U}' \subseteq \mathcal{X}'$ be quasi-compact open subsets over which f restricts to an isomorphism $\mathcal{U}' \simeq \mathcal{U}$. If Z and Z' are the respective set-theoretic complements of \mathcal{U} and \mathcal{U}' , then the induced square of pro-spectra*

$$\begin{array}{ccc} \{K^{\mathbb{B}}(\mathcal{X})\} & \longrightarrow & \widehat{K^{\mathbb{B}}}(\mathcal{X}'_{Z'}) \\ \downarrow & & \downarrow \\ \{K^{\mathbb{B}}(\mathcal{X}')\} & \longrightarrow & \widehat{K^{\mathbb{B}}}(\mathcal{X}'_{Z'}) \end{array}$$

is cartesian.

Remark 2.19. For *regular* (nonsingular) schemes and stacks, we will see in Subsect. 4.2 that the analogues of Theorems 2.17 and 2.18 do hold without passing to formal completions; see Theorems 4.10 and 4.11.

Remark 2.20. Alternatively, if we restrict our attention to affine (but possibly singular) derived schemes and only look at “very low” K-groups, then again Milnor excision and proper excision hold before formal completion; see [Mi, Thm 3.3], [B, Chap. XII, Thm. 8.3]. We will come back to this point in Subsect. 4.1.

2.6. Bass fundamental sequence. The Bass fundamental sequence is one of the few ways to understand negative K-groups. In fact, the construction of Bass–Thomason–Trobrough K-theory $K^{\mathbb{B}}$ is rigged to make it hold for the negative K-groups.

Theorem 2.21. *Let \mathcal{X} be a perfect derived Artin stack. Then for every integer n there is an exact sequence of abelian groups*

$$0 \rightarrow K_n(\mathcal{X}) \rightarrow K_n(\mathcal{X} \times \mathbf{A}^1) \oplus K_n(\mathcal{X} \times \mathbf{A}^1) \rightarrow K_n(\mathcal{X} \times \mathbf{G}_m) \xrightarrow{\partial} K_{n-1}(\mathcal{X}) \rightarrow 0,$$

functorial in \mathcal{X} with respect to inverse images. Moreover, the map ∂ admits a natural $K_(\mathcal{X})$ -module splitting.*

Proof. By Corollary 2.14 there is a Mayer–Vietoris sequence for the standard affine cover of $\mathcal{X} \times \mathbf{P}^1$:

$$\begin{aligned} \cdots \rightarrow K_{n+1}(\mathcal{X} \times \mathbf{G}_m) &\xrightarrow{\partial} K_n(\mathcal{X} \times \mathbf{P}^1) \rightarrow K_n(\mathcal{X} \times \mathbf{A}^1) \oplus K_n(\mathcal{X} \times \mathbf{A}^1) \\ &\rightarrow K_n(\mathcal{X} \times \mathbf{G}_m) \xrightarrow{\partial} \cdots \end{aligned}$$

Now apply the projective bundle formula (Theorem 2.10). The splitting comes from the Bott class $b \in K(\mathcal{X} \times \mathbf{G}_m)[-1]$. See [TT, Thm. 7.5] or [CiKh, Thm. 4.3.1, Rem. 4.3.2] for details. \square

2.7. Continuity.

Theorem 2.22. *Let \mathcal{X} be a perfect derived Artin stack. If \mathcal{X} is the limit of a cofiltered system $(\mathcal{X}_\alpha)_\alpha$ of perfect derived Artin stacks with affine transition morphisms, then the canonical map*

$$\varinjlim_{\alpha} \mathbf{K}^{\mathbf{B}}(\mathcal{X}_\alpha) \rightarrow \mathbf{K}^{\mathbf{B}}(\mathcal{X})$$

is invertible. In particular, there are canonical isomorphisms

$$\varinjlim_{\alpha} \mathbf{K}_n(X_\alpha) \simeq \mathbf{K}_n(X)$$

for all integers $n \in \mathbf{Z}$.

Proof. Combine Theorem 1.52(ii) and Theorem 2.5. □

2.8. Nil-invariance. The following result shows that, up to \mathbf{K}_{n+1} , the \mathbf{K} -groups of a derived commutative ring R are only sensitive to the first n homotopy groups of R .

Proposition 2.23. *Let R be a derived commutative ring. For every integer n , consider the canonical homomorphism $R \rightarrow \tau_{\leq n}(R)$ to the n -truncation (set $\tau_{\leq n}(R) := \tau_{\leq 0}(R) = \pi_0(R)$ for $n < 0$). Then the inverse image map*

$$\mathbf{K}^{\mathbf{B}}(\mathrm{Spec}(R)) \rightarrow \mathbf{K}^{\mathbf{B}}(\mathrm{Spec}(\tau_{\leq k}R))$$

induces an isomorphism $\mathbf{K}_i(\mathrm{Spec}(R)) \simeq \mathbf{K}_i(\mathrm{Spec}(\tau_{\leq k}R))$ for all $i \leq k + 1$.

Proof. See [Kh2, Prop. 4.2] or [BKRS, Prop. 5.1.3]. □

In particular, an affine derived scheme X and its classical truncation X_{cl} have the same \mathbf{K} -groups up to \mathbf{K}_1 .

Proposition 2.24 (Nil-invariance). *Let $i : Z \rightarrow X$ be a surjective closed immersion of affine derived schemes. Then the map $i^* : \mathbf{K}^{\mathbf{B}}(X) \rightarrow \mathbf{K}^{\mathbf{B}}(Z)$ is 1-connective; i.e., it is surjective on \mathbf{K}_1 and bijective on \mathbf{K}_n for all $n \leq 0$.*

Proof. By Proposition 2.23 we may replace X and Z by their classical truncations. By the Bass fundamental sequence (Theorem 2.21) it suffices to show surjectivity on \mathbf{K}_1 and bijectivity on \mathbf{K}_0 . This is [B, Chap. IX, Prop. 1.3]. □

Remark 2.25. Proposition 2.24 does not generalize to higher \mathbf{K} -groups. Contrast with Theorem 4.9.

Remark 2.26. For non-affine derived schemes or algebraic spaces, Proposition 2.24 still holds below the Krull dimension d (i.e., any surjective closed immersion will induce a bijection on \mathbf{K}_n for $n < -d$). For derived Artin stacks (with affine diagonal and nice stabilizers), the same holds if we replace d by the Nisnevich cohomological dimension (see [BKRS, Cor. 5.1.4]).

3. G-THEORY

3.1. Definition and basic properties. For a noetherian derived Artin stack \mathcal{X} , the *G-theory spectrum* $G(\mathcal{X})$ is defined as the algebraic K-theory of the stable ∞ -category of coherent complexes on \mathcal{X} :

$$G(\mathcal{X}) = K(\mathbf{D}_{\text{coh}}(\mathcal{X})).$$

Proposition 3.1. *Let \mathcal{X} be a noetherian derived Artin stack. Then the canonical map*

$$G(\mathcal{X}) = K(\mathbf{D}_{\text{coh}}(\mathcal{X})) \rightarrow K^{\text{B}}(\mathbf{D}_{\text{coh}}(\mathcal{X}))$$

is invertible. In other words, the spectrum $K^{\text{B}}(\mathbf{D}_{\text{coh}}(\mathcal{X}))$ is connective.

Proof. Since $\mathbf{D}_{\text{coh}}(\mathcal{X})$ admits a bounded t-structure with noetherian heart, this follows from [AGH, Thm. 1.2]. \square

Remark 3.2. Most of our discussion on G-theory can be extended from noetherian to qcqs stacks without modification. Proposition 3.1 seems to be an exception: I don't know whether $K^{\text{B}}(\mathbf{D}_{\text{coh}}(\mathcal{X}))$ will be connective for \mathcal{X} non-noetherian (cf. [AGH, Conj. B]).

Proposition 3.3. *Let \mathcal{X} be a noetherian derived Artin stack. Denote by $\text{Coh}(\mathcal{X})$ the abelian category of coherent sheaves on \mathcal{X} and by $K(\text{Coh}(\mathcal{X}))$ its K-theory in the sense of Quillen. Then there is a canonical isomorphism*

$$K(\text{Coh}(\mathcal{X})) \rightarrow G(\mathcal{X}),$$

functorial in \mathcal{X} .

Proof. Since $\mathbf{D}_{\text{coh}}(\mathcal{X})$ admits a bounded t-structure with heart $\text{Coh}(\mathcal{X})$, this follows from [Ba1, Thm. 6.1]. \square

Corollary 3.4 (Derived invariance). *Let \mathcal{X} be a noetherian derived Artin stack and write $i : \mathcal{X}_{\text{cl}} \rightarrow \mathcal{X}$ for the inclusion of the classical truncation. Then the direct image map*

$$i_* : G(\mathcal{X}_{\text{cl}}) \rightarrow G(\mathcal{X})$$

is invertible.

Proof. Follows immediately from Proposition 3.3, since $i_* : \mathbf{D}_{\text{coh}}(\mathcal{X}_{\text{cl}}) \rightarrow \mathbf{D}_{\text{coh}}(\mathcal{X})$ is t-exact and induces an equivalence $\text{Coh}(\mathcal{X}_{\text{cl}}) \simeq \text{Coh}(\mathcal{X})$ on hearts. \square

Theorem 3.5 (Poincaré duality). *Let \mathcal{X} be a noetherian derived Artin stack. If \mathcal{X} has bounded structure sheaf, then the inclusion $\mathbf{D}_{\text{perf}}(\mathcal{X}) \subseteq \mathbf{D}_{\text{coh}}(\mathcal{X})$ induces canonical maps*

$$K(\mathcal{X}) \rightarrow K^{\text{B}}(\mathcal{X}) \rightarrow K^{\text{B}}(\mathbf{D}_{\text{coh}}(\mathcal{X})) \simeq G(\mathcal{X})$$

which are invertible if \mathcal{X} is regular.

Proof. If \mathcal{X} is regular, then the inclusion $\mathbf{D}_{\text{perf}}(\mathcal{X}) \subseteq \mathbf{D}_{\text{coh}}(\mathcal{X})$ is an equality, hence the second map is invertible. From this it follows that $\mathbf{K}^{\text{B}}(\mathcal{X})$ is connective, so the first map is also invertible (since it is a connective cover). \square

Remark 3.6. The map $\mathbf{K}(\mathcal{X}) \rightarrow \mathbf{G}(\mathcal{X})$, or its factorization through $\mathbf{K}^{\text{B}}(\mathcal{X})$, is sometimes called the *Cartan map*.

3.2. Operations. Since tensoring with a perfect complex preserves coherence, we get a cap product

$$\cap : \mathbf{K}^{\text{B}}(\mathcal{X}) \otimes \mathbf{G}(\mathcal{X}) \rightarrow \mathbf{G}(\mathcal{X})$$

which is part of a $\mathbf{K}^{\text{B}}(\mathcal{X})$ -module structure on $\mathbf{G}(\mathcal{X})$.

For any morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ of finite Tor-amplitude, the inverse image functor $f^* : \mathbf{D}_{\text{coh}}(\mathcal{Y}) \rightarrow \mathbf{D}_{\text{coh}}(\mathcal{X})$ gives rise to a Gysin map

$$f_* : \mathbf{G}(\mathcal{Y}) \rightarrow \mathbf{G}(\mathcal{X}).$$

This is compatible with the K-theoretic inverse image under the maps in Theorem 3.5.

Let f be a proper morphism. If f is representable, or more generally of finite cohomological dimension, then there is a direct image map

$$f_* : \mathbf{G}(\mathcal{X}) \rightarrow \mathbf{G}(\mathcal{Y}).$$

This is compatible with the K-theoretic direct image (when f is of finite Tor-amplitude and almost of finite presentation) under the maps in Theorem 3.5. By Theorem 1.20 it is also $\mathbf{K}^{\text{B}}(\mathcal{Y})$ -linear (a homomorphism of $\mathbf{K}^{\text{B}}(\mathcal{Y})$ -module spectra). In other words:

Proposition 3.7 (Projection formula). *If $f : \mathcal{X} \rightarrow \mathcal{Y}$ is a proper and of finite cohomological dimension morphism of noetherian derived Artin stacks, then we have canonical identifications*

$$y \cap f_*(x) \simeq f_*(f^*(y) \cap x)$$

in $\mathbf{G}(\mathcal{Y})$, for all $x \in \mathbf{G}(\mathcal{X})$, $y \in \mathbf{K}^{\text{B}}(\mathcal{Y})$.

Theorem 1.20 similarly implies:

Proposition 3.8 (Base change formula). *Suppose given a homotopy cartesian square of noetherian derived Artin stacks*

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{g} & \mathcal{Y}' \\ \downarrow p & & \downarrow q \\ \mathcal{X} & \xrightarrow{f} & \mathcal{Y}. \end{array}$$

If f is proper and of finite cohomological dimension and q is of finite Tor-amplitude, then we have a canonical homotopy

$$q^* f_* \simeq g_* p^*$$

of maps $\mathbf{G}(\mathcal{X}) \rightarrow \mathbf{G}(\mathcal{Y}')$.

For G-theory we have the following analogue of Theorem 2.9:

Theorem 3.9 (Localization). *Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a closed immersion of noetherian derived Artin stacks with complementary open immersion $j : \mathcal{U} \rightarrow \mathcal{X}$. Then there is a canonical exact triangle of spectra*

$$G(\mathcal{Z}) \xrightarrow{i_*} G(\mathcal{X}) \xrightarrow{j^*} G(\mathcal{U}).$$

Proof. By Proposition 3.3, this follows from Quillen's dévissage and localization theorems [Q2, Sect. 5, Thms. 4 and 5]. \square

Corollary 3.10 (Nil invariance). *Let $i : Z \rightarrow X$ be a surjective closed immersion of noetherian derived Artin stacks. Then the induced morphism*

$$i_* : G(\mathcal{Z}) \rightarrow G(\mathcal{X})$$

is invertible.

3.3. Excision.

Theorem 3.11 (Étale excision). *Let \mathcal{X} be a noetherian derived Artin stack. Let $j : \mathcal{U} \rightarrow \mathcal{X}$ be an open immersion, and $Z \subseteq |\mathcal{X}|$ the set-theoretic complement. Then for any étale neighbourhood $f : \mathcal{X}' \rightarrow \mathcal{X}$ of Z (i.e., f is an étale morphism which restricts to an isomorphism $f^{-1}(Z)_{\text{red}} \rightarrow Z_{\text{red}}$), the induced square*

$$\begin{array}{ccc} G(\mathcal{X}) & \longrightarrow & G(\mathcal{U}) \\ \downarrow & & \downarrow \\ G(\mathcal{X}') & \longrightarrow & G(f^{-1}(\mathcal{U})) \end{array}$$

is cartesian.

Proof. By Theorem 3.9, the horizontal fibres are isomorphic. \square

Theorem 3.12 (Proper co-excision). *Let \mathcal{X} be a noetherian derived Artin stack. Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a closed immersion with open complement $j : \mathcal{U} \rightarrow \mathcal{X}$. Then for any proper morphism $f : \mathcal{X}' \rightarrow \mathcal{X}$ which restricts to an isomorphism $f^{-1}(\mathcal{U})_{\text{red}} \rightarrow \mathcal{U}_{\text{red}}$, the induced square*

$$\begin{array}{ccc} G(f^{-1}(\mathcal{Z})) & \longrightarrow & G(\mathcal{X}') \\ \downarrow & & \downarrow f_* \\ G(\mathcal{Z}) & \xrightarrow{i_*} & G(\mathcal{X}) \end{array}$$

is cocartesian (and hence cartesian).

Proof. By Theorem 3.9, the horizontal cofibres are isomorphic. \square

3.4. Projective bundles and blow-ups. We have the following G-theory analogues of Theorems 2.10 and 2.11.

Theorem 3.13. *Let \mathcal{X} be a noetherian derived Artin stack, \mathcal{E} a locally free sheaf on \mathcal{X} of rank $n + 1$, $n \geq 0$, and $q : \mathbf{P}_{\mathcal{X}}(\mathcal{E}) \rightarrow \mathcal{X}$ the associated projective bundle. Then the maps*

$$G(X) \rightarrow G(\mathbf{P}_{\mathcal{X}}(\mathcal{E})), \quad x \mapsto q^*(x) \cup [\mathcal{O}(-k)]$$

induce an isomorphism

$$G(\mathbf{P}_{\mathcal{X}}(\mathcal{E})) \simeq \bigoplus_{k=0}^n G(\mathcal{X}).$$

Proof. The standard semi-orthogonal decomposition on $\mathbf{D}_{\text{qc}}(\mathbf{P}_{\mathcal{X}}(\mathcal{E}))$ (see [Kh4, Thm. 3.3]) restricts to coherent complexes, so the formula follows from Corollary 2.2. \square

Theorem 3.14. *Let \mathcal{X} be a noetherian derived Artin stack. For any quasi-smooth closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ of virtual codimension n , consider the blow-up square*

$$\begin{array}{ccc} \mathbf{P}_{\mathcal{Z}}(\mathcal{N}_{\mathcal{Z}/\mathcal{X}}) & \xrightarrow{i_D} & \text{Bl}_{\mathcal{Z}} \mathcal{X} \\ \downarrow q & & \downarrow p \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array} \quad (3.15)$$

Then the maps $p^* : G(\mathcal{X}) \rightarrow G(\text{Bl}_{\mathcal{Z}} \mathcal{X})$ and

$$G(\mathcal{Z}) \rightarrow G(\text{Bl}_{\mathcal{Z}} \mathcal{X}), \quad x \mapsto i_{D,*}(q^*(x) \cup [\mathcal{O}(-k)]),$$

induce an isomorphism

$$G(\text{Bl}_{\mathcal{Z}} \mathcal{X}) \simeq G(\mathcal{X}) \oplus \bigoplus_{k=1}^{n-1} G(\mathcal{Z}).$$

Proof. The standard semi-orthogonal decomposition on $\mathbf{D}_{\text{qc}}(\text{Bl}_{\mathcal{Z}} \mathcal{X})$ (see [Kh4, Thm. 4.3]) restricts to coherent complexes. Thus the formula follows from Corollary 2.2. \square

Corollary 3.16. *Let \mathcal{X} be a noetherian derived Artin stack. For any quasi-smooth closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$, the blow-up square (3.15) induces a cartesian square*

$$\begin{array}{ccc} G(\mathcal{X}) & \xrightarrow{i^*} & G(\mathcal{Z}) \\ \downarrow p^* & & \downarrow q^* \\ G(\text{Bl}_{\mathcal{Z}} \mathcal{X}) & \xrightarrow{i_D^*} & G(\mathbf{P}_{\mathcal{Z}}(\mathcal{N}_{\mathcal{Z}/\mathcal{X}})). \end{array}$$

Proof. Combine Theorems 3.13 and 3.14. \square

3.5. Homotopy invariance. Let \mathcal{X} be a noetherian derived Artin stack. Let \mathcal{E} be a locally free sheaf of finite rank, $\phi : \mathcal{E} \rightarrow \mathcal{O}_{\mathcal{X}}$ a surjective⁴ $\mathcal{O}_{\mathcal{X}}$ -module homomorphism, and \mathcal{F} its fibre (which is locally free of finite rank). The associated *affine bundle*

$$\pi : \mathbf{V}_{\mathcal{X}}(\mathcal{E}, \phi) \rightarrow \mathcal{X}$$

is the moduli of splittings of ϕ . It fits in a closed/open pair

$$\mathbf{P}_{\mathcal{X}}(\mathcal{F}^{\vee}) \hookrightarrow \mathbf{P}_{\mathcal{X}}(\mathcal{E}^{\vee}) \leftarrow \mathbf{V}_{\mathcal{X}}(\mathcal{E}, \phi)$$

and it is a torsor under the vector bundle $\mathbf{V}_{\mathcal{X}}(\mathcal{F}^{\vee}) \rightarrow \mathcal{X}$. Every vector bundle torsor arises via this construction, see e.g. the proof of [T1, Thm. 4.1].

Theorem 3.17. *Let \mathcal{X} be a noetherian derived Artin stack. For any (\mathcal{E}, ϕ) as above, let $\pi : \mathbf{V}_{\mathcal{X}}(\mathcal{E}, \phi) \rightarrow \mathcal{X}$ denote the associated affine bundle. Then the inverse image map*

$$\pi^* : \mathbf{G}(\mathcal{X}) \rightarrow \mathbf{G}(\mathbf{V}_{\mathcal{X}}(\mathcal{E}, \phi))$$

is invertible.

Proof. Note that π is smooth, so the map exists. Combine the localization triangle (Theorem 3.9)

$$\mathbf{G}(\mathbf{P}_{\mathcal{X}}(\mathcal{F}^{\vee})) \rightarrow \mathbf{G}(\mathbf{P}_{\mathcal{X}}(\mathcal{E}^{\vee})) \rightarrow \mathbf{G}(\mathbf{V}_{\mathcal{X}}(\mathcal{E}, \phi))$$

with the projective bundle formula (Theorem 3.13). \square

For any finite locally free sheaf \mathcal{E} , the affine bundle associated to the projection $\mathcal{E} \oplus \mathcal{O}_{\mathcal{X}} \rightarrow \mathcal{O}_{\mathcal{X}}$ is the vector bundle $\mathbf{V}_{\mathcal{X}}(\mathcal{E}^{\vee})$ classifying sections $\mathcal{O}_{\mathcal{X}} \rightarrow \mathcal{E}$. Therefore we in particular get homotopy invariance for vector bundles.

Corollary 3.18. *Let \mathcal{X} be a noetherian derived Artin stack. For any finite locally free sheaf \mathcal{E} on \mathcal{X} , let $\pi : \mathbf{V}_{\mathcal{X}}(\mathcal{E}) \rightarrow \mathcal{X}$ denote the vector bundle parametrizing cosections of \mathcal{E} . Then the inverse image map*

$$\pi^* : \mathbf{G}(\mathcal{X}) \rightarrow \mathbf{G}(\mathbf{V}_{\mathcal{X}}(\mathcal{E}))$$

is invertible.

We can also generalize this further to vector bundle stacks.

Corollary 3.19. *Let \mathcal{X} be a noetherian derived Artin stack. Assume that \mathcal{X} has affine stabilizers. For any perfect complex \mathcal{E} on \mathcal{X} of Tor-amplitude ≤ 0 , let $\pi : \mathbf{V}_{\mathcal{X}}(\mathcal{E}) \rightarrow \mathcal{X}$ denote the vector bundle stack parametrizing cosections of \mathcal{E} . Then the inverse image map*

$$\pi^* : \mathbf{G}(\mathcal{X}) \rightarrow \mathbf{G}(\mathbf{V}_{\mathcal{X}}(\mathcal{E}))$$

is invertible.

Proof. If \mathcal{E} is of Tor-amplitude $[0, 0]$, then it is finite locally free and we are in the case of Corollary 3.18.

If \mathcal{E} is of Tor-amplitude $[-1, -1]$, then $E := \mathbf{V}_{\mathcal{X}}(\mathcal{E})$ is the classifying stack of the vector bundle $E[-1] := \mathbf{V}_{\mathcal{X}}(\mathcal{E}[1])$. In this case, the canonical morphism

⁴on π_0

$\sigma : X \twoheadrightarrow [X/E[-1]] \simeq E$ is the projection of a vector bundle (it is the quotient by $E[-1]$ of the projection $E[-1] \rightarrow X$). Thus σ^* is invertible by Corollary 3.18. Since $\sigma^*\pi^* = \text{id}$, we deduce that π^* is invertible. Repeating the same argument inductively also gives the case of Tor-amplitude $[-k, -k]$, for any $k \geq 0$.

Finally let \mathcal{E} be of Tor-amplitude $[-k, 0]$ for some $k > 0$. Since \mathcal{X} has affine stabilizers, its classical truncation admits a stratification by stacks with the resolution property (see [HR1, Prop. 2.6(i)]). By Corollary 3.4 and Theorem 3.9, we may therefore assume that \mathcal{X} admits the resolution property. In this case we can find a morphism $\phi : \mathcal{E}_0[-k] \rightarrow \mathcal{E}$ with \mathcal{E}_0 finite locally free, whose cofibre \mathcal{E}' is of Tor-amplitude $[-k+1, 0]$. Write $E := \mathbf{V}_{\mathcal{X}}(\mathcal{E})$, $E' := \mathbf{V}_{\mathcal{X}}(\mathcal{E}')$, and $E_0[k] := \mathbf{V}_{\mathcal{X}}(\mathcal{E}_0[-k])$. The projection $\pi : E \rightarrow \mathcal{X}$ factors through the morphism $\phi : E \rightarrow E_0[k]$ and the projection $\pi_0 : E_0[k] \rightarrow \mathcal{X}$. Since ϕ is the projection of a vector bundle (it is the quotient by $E_0[k-1]$ of the projection $\pi' : E' \rightarrow \mathcal{X}$), ϕ^* is invertible by the Tor-amplitude $[0, 0]$ case. By the $[-k, -k]$ case above, π_0^* is also invertible. Hence $\pi^* \simeq \phi^*\pi_0^*$ is invertible. \square

4. HOMOTOPY INVARIANCE AND SINGULARITIES

4.1. Weibel’s conjecture. Let X be a noetherian algebraic space. If X is regular, then combining Poincaré duality (Theorem 3.5) with Proposition 3.1 and Corollary 3.18 yields:

- (i) The negative K-groups $K_{-n}(X)$ vanish for all $n > 0$.
- (ii) For any vector bundle $\pi : E \rightarrow X$, the map $\pi^* : K^{\mathbf{B}}(X) \rightarrow K^{\mathbf{B}}(E)$ is invertible.

When X is singular, both these properties fail. In fact, as we will see in the next subsection, all the “pathological” behaviour of algebraic K-theory for singular schemes, such as failure of nil-invariance and excision of open subsets (see Subsects. 2.8 and 2.5) can be traced back to the failure of homotopy invariance. In particular, these excision statements will turn out to be true for nonsingular spaces.

On the other hand, many of these pathologies disappear on “low enough” K-groups (see Proposition 2.24 and Remark 2.20). This is in some sense “explained” by the following statement, which says that even for singular spaces, homotopy invariance does hold on low enough K-groups.

Theorem 4.1. *Let X be a noetherian algebraic space. Suppose that X is pro-smooth over a noetherian algebraic space of Krull dimension d . Then we have:*

- (i) *The negative K-groups $K_{-n}(X)$ vanish for all $n > d$.*
- (ii) *For any vector bundle $\pi : E \rightarrow X$, the maps $\pi^* : K_{-n}(X) \rightarrow K_{-n}(E)$ are invertible for all $n \geq d$.*

Example 4.2. If X is smooth (or pro-smooth) over a field, then it is regular and Theorem 4.1 just recovers the nonsingular situation discussed above. (Conversely, if we restrict our attention to affine schemes over a field, then by Popescu’s desingularization theorem [Spi, Thm. 1.1], regularity implies pro-smoothness.)

Example 4.3. For X a singular scheme of dimension d , Theorem 4.1 was known as Weibel’s conjecture (formulated in [We1, Questions 2.9] for X affine). Weibel’s conjecture was proven by Kerz–Strunk–Tamme [KST, Thm. B] for schemes.

Remark 4.4. Theorem 4.1 also holds for Deligne–Mumford stacks, or more generally Artin stacks with quasi-finite diagonal; see [BKRS, Thm. D, Rem. 5.3.3]. In fact, there is also a statement for algebraic stacks with affine diagonal and nice stabilizers, but Krull dimension has to be replaced by a weaker bound.

Proof of Theorem 4.1. By assumption, there exists a noetherian algebraic space S of dimension d and a cofiltered system of smooth S -schemes $(X_\alpha)_\alpha$ with affine transition maps, whose limit is X . By continuity (Theorem 2.22) there are canonical isomorphisms

$$K_{-n}(X) \simeq \varinjlim_{\alpha} K_{-n}(X_\alpha)$$

for all $n \in \mathbf{Z}$. Hence (i) would follow from the analogous claims for X_α . Similarly for (ii), since by Corollary 1.54 the vector bundle $\pi : E \rightarrow X$ descends to X_α for every α .

Thus we may assume that X is smooth over S . In this case, the statement is proven by a slight variant on the proof of Weibel’s conjecture. See [BKRS, Rem. 5.3.3], or [Sa]⁵ in the case of schemes. \square

4.2. Homotopy invariant K-theory. We can introduce a variant of K-theory by *forcing* homotopy invariance for all (possibly singular) spaces. This construction goes back to Weibel [We2]. See [KrRa, HoKr, Ho4] for the case of stacks, [Kh4] for the case of derived algebraic spaces, and [KhRa] for derived stacks. We restrict to algebraic spaces for simplicity of exposition.

Let Δ^\bullet denote the standard cosimplicial affine scheme whose n th term is affine n -space \mathbf{A}^n (see e.g. [MV, p. 45]). For any qcqs derived algebraic space X , let $\mathrm{KH}(X)$ denote the geometric realization of the simplicial diagram $K^{\mathrm{B}}(\Delta^\bullet)$:

$$\mathrm{KH}(X) = \varinjlim_{[n] \in \Delta^{\mathrm{op}}} K^{\mathrm{B}}(X \times \Delta^n).$$

The canonical map of presheaves $K^{\mathrm{B}} \rightarrow \mathrm{KH}$ exhibits KH as the \mathbf{A}^1 -localization of K-theory in the sense of \mathbf{A}^1 -homotopy theory (see [Ci]).

⁵Thanks to the referee for pointing out this reference.

Remark 4.5. If X has bounded structure sheaf then by the universal property of \mathbf{A}^1 -localization and homotopy invariance for \mathbf{G} -theory (Corollary 3.18), we find that the canonical map $\mathbf{K}^{\mathbf{B}}(X) \rightarrow \mathbf{G}(X)$ (Theorem 3.5) factors through a map

$$\mathrm{KH}(X) \rightarrow \mathbf{G}(X),$$

functorial with respect to finite Tor-amplitude inverse images. If X is regular, then so are $X \times \Delta^n$ for all n , so this map is invertible. Hence all the maps

$$\mathbf{K}(X) \rightarrow \mathbf{K}^{\mathbf{B}}(X) \rightarrow \mathrm{KH}(X) \rightarrow \mathbf{G}(X)$$

are invertible when X is regular.

Remark 4.6. Since colimits of spectra are exact, KH inherits the following properties from $\mathbf{K}^{\mathbf{B}}$: localization sequence (Theorem 2.9), excision of closed subsets (Theorem 2.13), projective bundle formula (Theorem 2.10), and blow-up formula (Theorem 2.11).

Remark 4.7. For any \mathbf{Z} -linear stable ∞ -category \mathcal{A} we can define $\mathrm{KH}(\mathcal{A})$ as the spectrum

$$\mathrm{KH}(\mathcal{A}) = \varinjlim_{n \in \Delta^{\mathrm{op}}} \mathbf{K}^{\mathbf{B}}(\mathcal{A} \otimes \mathbf{D}_{\mathrm{perf}}(\Delta^n))$$

so that $\mathrm{KH}(X) = \mathrm{KH}(\mathbf{D}_{\mathrm{perf}}(X))$. Therefore KH also has cup products and direct images along proper morphisms that are almost of finite presentation and of finite Tor-amplitude. These satisfy the projection and base change formulas (Propositions 2.7 and 2.8).

4.3. Properties of KH . As hinted at earlier, passing from $\mathbf{K}^{\mathbf{B}}$ to KH has the remarkable side effect of forcing several other properties that were only true in \mathbf{K} -theory either for regular (nonsingular) stacks, or only for very low K -groups for affines.

Proposition 4.8 (Homotopy invariance). *For any qcqs derived algebraic space X and any vector bundle $\pi : E \rightarrow X$, the map $\pi^* : \mathrm{KH}(X) \rightarrow \mathrm{KH}(E)$ is invertible.*

Proof. Using Theorem 2.13(ii), we may assume the bundle is trivial. By induction we may assume it is of rank one. In that case this holds by construction. \square

Theorem 4.9 (Nil-invariance). *Let $i : Z \rightarrow X$ be a surjective closed immersion of qcqs derived algebraic spaces. Then the map $i^* : \mathrm{KH}(X) \rightarrow \mathrm{KH}(Z)$ is invertible.*

Proof. See [Kh4, Thm. 5.13]. \square

In particular, taking Z to be the classical truncation of X shows that KH is insensitive to derived structures.

We also have the following versions of “excising open subsets”.

Theorem 4.10 (Milnor excision). *For any Milnor square of qcqs derived algebraic spaces as in Subsect. 2.5, the induced square*

$$\begin{array}{ccc} \mathrm{KH}(X) & \xrightarrow{i^*} & \mathrm{KH}(Z) \\ \downarrow f^* & & \downarrow \\ \mathrm{KH}(X') & \longrightarrow & \mathrm{KH}(Z') \end{array}$$

is cartesian.

Proof. By Theorem 4.9 we may replace the square by its classical truncation. By Theorem 2.13(ii) and [Lu4, Thm. 3.4.2.1] we may assume it consists of affine schemes. In this case the claim is a classical result of Weibel (it is equivalent to [We2, Thm. 2.1]). \square

Theorem 4.11 (Proper excision). *Let $f : X' \rightarrow X$ be a proper morphism of qcqs derived algebraic spaces. Let $U \subseteq X$ and $U' \subseteq X'$ be quasi-compact open subsets such that f restricts to an isomorphism $U'_{\mathrm{red}} \rightarrow U_{\mathrm{red}}$, and let $Z \subseteq X$ and $Z' \subseteq X'$ be their reduced complements. Then the square*

$$\begin{array}{ccc} \mathrm{KH}(X) & \longrightarrow & \mathrm{KH}(Z) \\ \downarrow f^* & & \downarrow \\ \mathrm{KH}(X') & \longrightarrow & \mathrm{KH}(Z') \end{array}$$

is cartesian.

Proof. For schemes this is the main result of [Ci] (where the noetherian assumption can be dropped by [Ho1, App. C]). For the case of algebraic spaces see [Kh4, Thm. D]. \square

Finally we also get the following Mayer–Vietoris property for closed subspaces.

Corollary 4.12. *Let X be a qcqs derived algebraic space. For any closed subspaces Y and Z such that $X = Y \cup Z$, the square*

$$\begin{array}{ccc} \mathrm{KH}(X) & \longrightarrow & \mathrm{KH}(Y) \\ \downarrow & & \downarrow \\ \mathrm{KH}(Z) & \longrightarrow & \mathrm{KH}(Y \cap Z) \end{array}$$

is cartesian. Here $Y \cap Z$ is the classical or derived scheme-theoretic intersection (KH is insensitive to the difference by Theorem 4.9).

5. RATIONAL ÉTALE K-THEORY AND G-THEORY

Given a derived Artin stack \mathcal{X} , let $\mathrm{K}^{\mathrm{B}}(\mathcal{X})_{\mathbf{Q}}$ and $\mathrm{G}(\mathcal{X})_{\mathbf{Q}}$ denote the rationalized K- and G-theory spectra, respectively. In this section we introduce étale-localized versions $\mathrm{K}^{\mathrm{ét}}(\mathcal{X})_{\mathbf{Q}}$ and $\mathrm{G}^{\mathrm{ét}}(\mathcal{X})_{\mathbf{Q}}$.

5.1. Descent on algebraic spaces.

Theorem 5.1 (Étale and flat descent).

- (i) *On the ∞ -category of qcqs derived algebraic spaces, the presheaf $X \mapsto \mathbf{K}^{\mathbf{B}}(X)_{\mathbf{Q}}$ satisfies descent for the étale and finite flat topologies.*
- (ii) *On the ∞ -category of qcqs derived algebraic spaces, the presheaf $X \mapsto \mathbf{K}\mathbf{H}(X)_{\mathbf{Q}}$ satisfies descent for the fppf topology.*
- (iii) *On the ∞ -category of noetherian derived algebraic spaces and finite Tor-amplitude morphisms, the presheaf $X \mapsto \mathbf{G}(X)_{\mathbf{Q}}$ satisfies descent for the fppf topology.*

Remark 5.2. Recall that any smooth surjection generates a covering for the étale topology. Thus in particular, Theorem 5.1 implies that for any smooth surjection $p : U \rightarrow X$, the inverse image map p^* induces an isomorphism between $\mathbf{K}^{\mathbf{B}}(X)_{\mathbf{Q}}$ and the homotopy limit of the cosimplicial diagram

$$\mathbf{K}^{\mathbf{B}}(U)_{\mathbf{Q}} \rightrightarrows \mathbf{K}^{\mathbf{B}}(U \times_X U)_{\mathbf{Q}} \rightrightarrows \mathbf{K}^{\mathbf{B}}(U \times_X U \times_X U)_{\mathbf{Q}} \rightrightarrows \dots$$

and similarly for KH and G-theory.

Proof of Theorem 5.1. We first show that each presheaf satisfies finite flat descent. For this we may restrict to the small étale site of a fixed derived algebraic space S . Combining Theorem 2.13(ii) or its KH/G-theory analogue (Remark 4.6 or Corollary 3.11) with [Lu4, Thm. 3.4.2.1], we may assume that S is affine. Then the claim follows by applying [CMNN, Prop. 5.4] (taking $n = 0$, cf. Appendix A in *op. cit.*), which is a statement about localizing invariants of $\mathbf{D}_{\text{perf}}(S)$ -linear ∞ -categories (see Def. A.1 of *op. cit.*) that take values in rational spectra. For the K-theory statement, take the localizing invariant that sends a $\mathbf{D}_{\text{perf}}(S)$ -linear ∞ -category \mathcal{A} to the spectrum $\mathbf{K}^{\mathbf{B}}(\mathcal{A})_{\mathbf{Q}}$. For KH, take the localizing invariant

$$\mathcal{A} \mapsto \varinjlim_{[n] \in \Delta^{\text{op}}} \mathbf{K}^{\mathbf{B}}(\mathcal{A} \otimes_{\mathbf{D}_{\text{perf}}(S)} \mathbf{D}_{\text{perf}}(S \times \Delta^n))_{\mathbf{Q}}.$$

For G-theory, take the localizing invariant K' that sends a $\mathbf{D}_{\text{perf}}(S)$ -linear ∞ -category \mathcal{A} to the spectrum $\mathbf{K}(\mathcal{A} \otimes_{\mathbf{D}_{\text{perf}}(S)} \text{Coh}(S))_{\mathbf{Q}}$. Indeed, for every X étale over S , there is a canonical isomorphism

$$\begin{aligned} K'(\mathbf{D}_{\text{perf}}(X)) &= \mathbf{K}(\mathbf{D}_{\text{perf}}(X) \otimes_{\mathbf{D}_{\text{perf}}(S)} \text{Coh}(S))_{\mathbf{Q}} \\ &\simeq \mathbf{K}(\text{Coh}(X))_{\mathbf{Q}} = \mathbf{G}(X)_{\mathbf{Q}} \end{aligned}$$

by [GR, Chap. 4, Rem. 3.3.3].

By [Lu4, Thm. B.6.4.1], combining finite flat descent with Nisnevich descent (Remark 2.15) yields étale descent for each presheaf.

For $\mathbf{K}\mathbf{H}_{\mathbf{Q}}$ and $\mathbf{G}_{\mathbf{Q}}$, we may use derived invariance (Theorem 4.9 and Corollary 3.4) to restrict to the site of classical algebraic spaces. Since the fppf topology on this site is generated by Nisnevich coverings and finite flat coverings (see e.g. [SP, Tag 05WN]), the claim follows. \square

Theorem 5.3 (Proper descent). *On the ∞ -category of qcqs derived algebraic spaces, the presheaf $X \mapsto \mathrm{KH}(X)_{\mathbf{Q}}$ satisfies descent along proper schematic surjections of finite presentation. That is, for any proper schematic surjection of finite presentation $p : Y \rightarrow X$, p^* induces an isomorphism between $\mathrm{KH}(X)_{\mathbf{Q}}$ and the homotopy limit of the cosimplicial diagram*

$$\mathrm{KH}(Y)_{\mathbf{Q}} \rightrightarrows \mathrm{KH}(Y \times_X^{\mathbf{R}} Y)_{\mathbf{Q}} \rightrightarrows \mathrm{KH}(Y \times_X^{\mathbf{R}} Y \times_X^{\mathbf{R}} Y)_{\mathbf{Q}} \rightrightarrows \dots$$

Proof. By derived invariance (Theorem 4.9) and descent (Corollary 5.1), we may restrict to the site of classical schemes. Furthermore we may fix a scheme S and restrict to the site of S -schemes of finite presentation. Then we may apply the criterion of [BS, Thm. 2.9], which reduces the claim to fppf descent (Theorem 5.1) and proper cdh descent (Theorem 4.11). \square

Remark 5.4. Presumably, Theorem 5.3 extends to (non-schematic) proper surjections of finite presentation, but I did not check that [BS, Thm. 2.9] extends to algebraic spaces.

Remark 5.5. In fact, the previous proof shows that $\mathrm{KH}_{\mathbf{Q}}$ satisfies descent for Voevodsky’s h-topology (or rather, Rydh’s non-noetherian generalization of it [Ry1, Sect. 8]) on the site of (derived) schemes.

5.2. Étale K-theory and G-theory. Theorem 5.1 does not extend to stacks. In fact, on derived Artin stacks, rational G-theory only satisfies descent with respect to étale maps that are *isovariant* or “stabilizer-preserving” (see [Jo, Sect. 3]).

Nevertheless, Theorem 5.1 implies that $\mathrm{K}^{\mathbf{B}}(-)_{\mathbf{Q}}$, $\mathrm{KH}(-)_{\mathbf{Q}}$, and $\mathrm{G}(-)_{\mathbf{Q}}$ extend uniquely to étale sheaves on derived Artin stacks, which we denote by

$$\mathrm{K}^{\text{ét}}(-)_{\mathbf{Q}}, \mathrm{KH}^{\text{ét}}(-)_{\mathbf{Q}}, \text{ and } \mathrm{G}^{\text{ét}}(-)_{\mathbf{Q}}.$$

These could equivalently be described as the étale localizations of the presheaves $\mathrm{K}^{\mathbf{B}}(-)_{\mathbf{Q}}$, $\mathrm{KH}(-)_{\mathbf{Q}}$, and $\mathrm{G}(-)_{\mathbf{Q}}$, respectively. Note that, as before, we only ever discuss descent with respect to Čech covers and not hypercovers.

Thus by construction, for any derived Artin stack \mathcal{X} and any smooth atlas $p : X \rightarrow \mathcal{X}$, $\mathrm{K}^{\text{ét}}(\mathcal{X})_{\mathbf{Q}}$ is isomorphic to the homotopy limit of the cosimplicial diagram

$$\mathrm{K}^{\text{ét}}(X)_{\mathbf{Q}} \rightrightarrows \mathrm{K}^{\mathbf{B}}(X \times_{\mathcal{X}} X)_{\mathbf{Q}} \rightrightarrows \mathrm{K}^{\mathbf{B}}(X \times_{\mathcal{X}} X \times_{\mathcal{X}} X)_{\mathbf{Q}} \rightrightarrows \dots$$

and similarly for KH and G-theory. For example, if $\mathcal{X} = [X/G]$ is the quotient of a derived algebraic space X by the action of an fppf group space G , then $\mathrm{K}^{\text{ét}}([X/G])_{\mathbf{Q}}$ is the Borel-type G -equivariant K-theory of X :

$$\mathrm{K}^{\text{ét}}(X)_{\mathbf{Q}} \rightrightarrows \mathrm{K}^{\mathbf{B}}(G \times X)_{\mathbf{Q}} \rightrightarrows \mathrm{K}^{\mathbf{B}}(G \times G \times X)_{\mathbf{Q}} \rightrightarrows \dots$$

Similarly for $\mathrm{KH}^{\text{ét}}(-)_{\mathbf{Q}}$ and $\mathrm{G}^{\text{ét}}(-)_{\mathbf{Q}}$.

All properties of K-theory, G-theory, and KH-theory involving *inverse* image functoriality easily extend to the étale-local variants just by descent from the case of algebraic spaces. This applies for example to the projective bundle

formula, excision for quasi-smooth blow-ups, and homotopy invariance. In particular, we can drop the extra hypothesis in the latter statement (compare with Corollary 3.19).

Corollary 5.6 (Homotopy invariance). *Let \mathcal{X} be a derived Artin stack. Let \mathcal{E} be a perfect complex on \mathcal{X} of Tor-amplitude ≤ 0 , and $\pi : \mathbf{V}_{\mathcal{X}}(\mathcal{E}) \rightarrow \mathcal{X}$ the associated vector bundle stack. Then inverse image along π induces isomorphisms of spectra*

$$\begin{aligned} \pi^* : \mathrm{KH}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}} &\rightarrow \mathrm{KH}^{\acute{e}t}(\mathbf{V}_{\mathcal{X}}(\mathcal{E}))_{\mathbf{Q}}, \\ \pi^* : \mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}} &\rightarrow \mathrm{G}^{\acute{e}t}(\mathbf{V}_{\mathcal{X}}(\mathcal{E}))_{\mathbf{Q}}. \end{aligned}$$

Proof. One argues as in Corollary 3.19, except that the resolution property can be guaranteed simply by localizing on \mathcal{X} (in the étale topology). \square

In contrast, the direct image functoriality is somewhat subtle (see further discussion in Subsect. 5.4 below). For any proper *representable* morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ between noetherian derived Artin stacks, there is indeed a direct image map

$$f_* : \mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}} \rightarrow \mathrm{G}^{\acute{e}t}(\mathcal{Y})_{\mathbf{Q}}.$$

Since properness is local on the target, descent allows us to reduce to the case where $\mathcal{Y} = Y$ is representable by a derived algebraic space. Since f is representable, it follows that $\mathcal{X} = X$ is also an algebraic space. Thus we simply take the map

$$\mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}} \simeq \mathrm{G}(X)_{\mathbf{Q}} \xrightarrow{f_*} \mathrm{G}(Y)_{\mathbf{Q}} \simeq \mathrm{G}^{\acute{e}t}(\mathcal{Y})_{\mathbf{Q}}.$$

Properties of G-theory involving proper representable direct images also extend to the étale-local variants (e.g., localization, base change, projection formulas).

5.3. Proper codescent. Just as étale excision (Corollary 3.11) in G-theory extends to étale descent in rational G-theory (Theorem 5.1), proper co-excision (Theorem 3.12) similarly extends to proper co-descent with rational coefficients. This extends moreover to rational étale G-theory of stacks.

Theorem 5.7 (Proper codescent). *On the ∞ -category of noetherian derived Artin stacks, the copresheaf $\mathcal{X} \mapsto \mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}}$ satisfies descent along proper representable surjections.*

Remark 5.8. More concretely, Theorem 5.7 asserts that for any proper representable surjection $f : \mathcal{Z} \rightarrow \mathcal{X}$, f_* induces an isomorphism between $\mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}}$ and the homotopy colimit of the simplicial diagram

$$\cdots \rightrightarrows \mathrm{G}^{\acute{e}t}(\mathcal{Z} \times_{\mathcal{X}}^{\mathbf{R}} \mathcal{Z} \times_{\mathcal{X}}^{\mathbf{R}} \mathcal{Z})_{\mathbf{Q}} \rightrightarrows \mathrm{G}^{\acute{e}t}(\mathcal{Z} \times_{\mathcal{X}}^{\mathbf{R}} \mathcal{Z})_{\mathbf{Q}} \rightrightarrows \mathrm{G}^{\acute{e}t}(\mathcal{Z})_{\mathbf{Q}}.$$

If \mathcal{X} and \mathcal{Z} are classical, then the fibred products need not be derived (see Corollary 3.4).

We will need a few preliminary observations. The first is in fact a straightforward corollary of Theorem 5.7 (consider the diagonal, which is a surjective closed immersion), but we can also give a direct proof.

Proposition 5.9 (Topological invariance). *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a finite radicial surjection of noetherian derived Artin stacks (i.e., a universal homeomorphism of finite type). Then the induced morphism*

$$f_* : G^{\text{ét}}(\mathcal{X})_{\mathbf{Q}} \rightarrow G^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}}$$

is invertible.

Proof. By derived invariance (Corollary 3.4), we may assume \mathcal{X} and \mathcal{Y} are classical. By étale descent (and base change, Proposition 3.8) we may also assume $\mathcal{Y} = Y$ (and hence $\mathcal{X} = X$) is affine. By [EGA, IV₄, 17.16.4], there is a partition of Y by finitely many locally closed subschemes Y_i along with finite flat surjections $Y'_i \twoheadrightarrow Y_i$ such that each base change $f_i : X \times_Y Y'_i \rightarrow Y'_i$ admits a section. By localization (Theorem 3.9), we may replace Y by Y_i and by finite flat descent (Theorem 5.1) we may further replace Y_i by Y'_i . In other words, we may assume that f admits a section s . Since f is radicial, s is a surjective closed immersion. Then s_* is an isomorphism (Corollary 3.10), hence so is f_* . \square

Proof of Theorem 5.7. Let $f : \mathcal{Z} \rightarrow \mathcal{X}$ be a proper representable surjection. By nil-invariance (Corollary 3.10), we may assume \mathcal{X} and \mathcal{Z} are reduced classical stacks. By étale descent (and base change, Proposition 3.8) we may assume $\mathcal{X} = X$ is affine (and hence $\mathcal{Z} = Z$ is an algebraic space). By noetherian induction and localization (Theorem 3.9), we may further reduce to the case where X is the spectrum of a field k . Let z be a closed point of the (nonempty, noetherian) scheme Z . Its residue field $\kappa(z)$ is purely inseparable over a Galois extension k' of k . Finite étale descent (Theorem 5.1) in the case of $\text{Spec}(k') \rightarrow \text{Spec}(k)$ gives an isomorphism

$$G(k')^G_{\mathbf{Q}} \simeq G(k),$$

where the left-hand side is the spectrum of homotopy invariants with respect to the Galois group G . Similarly for the induced Galois cover $Z' = Z \times_X \text{Spec}(k') \rightarrow Z$. Thus the diagram

$$\cdots \rightrightarrows G^{\text{ét}}(Z \times_X^{\mathbf{R}} Z \times_X^{\mathbf{R}} Z)_{\mathbf{Q}} \rightrightarrows G(Z \times_X^{\mathbf{R}} Z)_{\mathbf{Q}} \rightrightarrows G(Z)_{\mathbf{Q}} \rightarrow G(X)_{\mathbf{Q}}$$

is identified with the colimit of the same diagram where Z and $X = \text{Spec}(k)$ are replaced by Z' and $X' := \text{Spec}(k')$. Since formation of homotopy invariants commutes with colimits (recall that with rational coefficients, the former coincide with homotopy coinvariants, which are colimits), it will suffice to show the claim after this replacement. By Proposition 5.9 we may also base change further along the finite radicial surjection $\text{Spec}(\kappa(z)) \rightarrow \text{Spec}(k')$. In particular, we may assume that $f : Z \rightarrow X$ admits a section s . In that case the claim is obvious, as s induces a splitting of the augmented simplicial object in question (which is then automatically a homotopy colimit diagram, see e.g. [Lu1, 6.1.3.16]). \square

5.4. Non-representable direct image. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper morphism between noetherian derived Artin stacks. We can use proper codescent to construct a direct image map

$$f_* : G^{\text{ét}}(\mathcal{X})_{\mathbf{Q}} \rightarrow G^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}}$$

even if f is not representable.

By [Ol, Thm. 1.1] (applied to the classical truncations), there exists a scheme Z and a proper surjection $g : Z \rightarrow \mathcal{X}$ such that the composite $Z \rightarrow \mathcal{X} \rightarrow \mathcal{Y}$ is projective (in particular, proper representable). By proper codescent along g (Theorem 5.7), we get an essentially unique map $f_* : G^{\text{ét}}(\mathcal{X})_{\mathbf{Q}} \rightarrow G^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}}$ extending the direct image maps

$$G^{\text{ét}}(Z \times_{\mathcal{X}} \cdots \times_{\mathcal{X}} Z)_{\mathbf{Q}} \rightarrow G^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}}.$$

Given another choice of Z and g , it is an easy exercise to show that the two morphisms f_* are homotopic. Similarly, if $f : \mathcal{X} \rightarrow \mathcal{Y}$ and $f' : \mathcal{Y} \rightarrow \mathcal{Z}$ are two proper morphisms (where \mathcal{X} and \mathcal{Y} both have quasi-finite separated diagonal), then the two maps $f'_* \circ f_*$ and $(f' \circ f)_*$ are homotopic. (We do not address coherence of these homotopies here.) These direct images also satisfy the base change and projection formulas (Propositions 3.7 and 3.8).

Warning 5.10. These non-representable direct images are *not* compatible with those in “genuine” G-theory induced by proper push-forward of coherent complexes. That is, let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper morphism of finite cohomological dimension and assume \mathcal{X} has quasi-finite separated diagonal. Then the square

$$\begin{array}{ccc} G(\mathcal{X})_{\mathbf{Q}} & \xrightarrow{f_*} & G(\mathcal{Y})_{\mathbf{Q}} \\ \downarrow & & \downarrow \\ G^{\text{ét}}(\mathcal{X})_{\mathbf{Q}} & \xrightarrow{f_*} & G^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}} \end{array}$$

typically does not commute (unless f is representable).

Consider the following standard example. Let k be a field, G a finite group, and $f : BG \rightarrow \text{Spec}(k)$ the structural map of the classifying stack. If the order of G is prime to the characteristic of k , then f is of finite cohomological dimension. Let $g : \text{Spec}(k) \rightarrow BG$ denote the quotient morphism, which is finite étale. Using the base change and projection formulas, one finds that both composites g^*g_* and g_*g^* on $G(k)$ are multiplication by the order of G . In particular, g induces an isomorphism

$$\mathbf{Q} \simeq G^{\text{ét}}(k)_{\mathbf{Q}} \xrightarrow{g_*} G^{\text{ét}}(BG)_{\mathbf{Q}}.$$

By functoriality, it follows that $f_* : G^{\text{ét}}(BG)_{\mathbf{Q}} \rightarrow G^{\text{ét}}(k)_{\mathbf{Q}}$ is homotopic to a scalar multiple of g^* . Since the étale localization map $G(BG)_{\mathbf{Q}} \rightarrow G^{\text{ét}}(BG)_{\mathbf{Q}}$ commutes with g^* by definition, we see that the square commutes if and only if f_* and g^* are also homotopic at the level of $G(-)_{\mathbf{Q}}$ (up to multiplication by a scalar). But under the identifications of $G(BG)_{\mathbf{Q}}$ with the K-theory spectrum of G -equivariant k -vector spaces and $G(k)_{\mathbf{Q}}$ with the K-theory spectrum of k -vector spaces, the map g^* sends the class of a G -representation

V to the underlying vector space V while f_* sends it to the class of the invariant subspace V^G . No scalar multiplication identifies these (e.g. take V to be a nontrivial representation of dimension one so that $V^G = 0$).

6. VIRTUAL FUNDAMENTAL CLASSES

6.1. Fundamental classes in G-theory. Let \mathcal{X} be a regular (nonsingular) Artin stack. K-theoretic Poincaré duality (Theorem 3.5) can be reformulated as the assertion that there is a canonical class $[\mathcal{X}]_G \in G(\mathcal{X})$ such that cap product gives an isomorphism

$$[\mathcal{X}]_G \cap (-) : K^B(\mathcal{X}) \rightarrow G(\mathcal{X}).$$

Of course, $[\mathcal{X}]_G$ is just the class of the structure sheaf $[\mathcal{O}_{\mathcal{X}}]$.

This construction can be extended to (derived) stacks that are possibly *singular* but still quasi-smooth. First note that by Lemma 1.15 we have:

Corollary 6.1. *Let $\mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-smooth morphism of derived Artin stacks. If \mathcal{Y} has bounded structure sheaf, then so does \mathcal{X} .*

Construction 6.2 (G-theoretic fundamental class). Let \mathcal{X} be a quasi-smooth derived Artin stack over a regular noetherian Artin stack \mathcal{S} . Then \mathcal{X} is noetherian and its structure sheaf $\mathcal{O}_{\mathcal{X}}$ is coherent by Corollary 6.1 and Remark 1.8. In particular, there is a *G-theoretic fundamental class*

$$[\mathcal{X}]_G := [\mathcal{O}_{\mathcal{X}}] \in G(\mathcal{X}).$$

Remark 6.3. Under the canonical isomorphism $G(\mathcal{X}) \simeq G(\mathcal{X}_{\text{cl}})$ (Corollary 3.4), the fundamental class $[\mathcal{X}]_G$ corresponds uniquely to a class

$$[\mathcal{X}]_G^{\text{vir}} \in G(\mathcal{X}_{\text{cl}}),$$

which we call the G-theoretic *virtual* fundamental class. Explicitly, it can be described as the alternating sum

$$[\mathcal{X}]_G^{\text{vir}} = \sum_{i \leq 0} (-1)^i [\mathbf{H}^i(\mathcal{O}_{\mathcal{X}})],$$

which is finite by assumption, where $\mathbf{H}^i(\mathcal{O}_{\mathcal{X}}) = \pi_{-i}(\mathcal{O}_{\mathcal{X}})$ are viewed as coherent sheaves on \mathcal{X}_{cl} via the equivalence $\text{Coh}(\mathcal{X}_{\text{cl}}) \simeq \mathbf{D}_{\text{coh}}(\mathcal{X})^\heartsuit$.

This virtual G-theory class agrees with the *virtual structure sheaf* studied in detail by Y.-P. Lee [Lee]. Construction 6.2 goes back to [Ko, 1.4.2] and was written down more precisely by Toën in [To3, §4.4, para. 3]. Note that the various properties of virtual structure sheaves [Lee, Subsect. 2.4] are completely transparent from Construction 6.2.

Remark 6.4. Since the structural morphism $f : \mathcal{X} \rightarrow \mathcal{S}$ is quasi-smooth and hence of finite Tor-amplitude (Lemma 1.15), there is an inverse image map

$$f^* : G(\mathcal{S}) \rightarrow G(\mathcal{X}).$$

and the fundamental class $[\mathcal{X}]_G$ is of course the same as the image of the fundamental class $[\mathcal{O}_{\mathcal{S}}] \in G(\mathcal{S})$ defined above.

Remark 6.5. More generally, for any quasi-smooth morphism of noetherian derived Artin stacks $f : \mathcal{X} \rightarrow \mathcal{Y}$ there is again by Lemma 1.15 a map

$$f^* : \mathbf{G}(\mathcal{Y}) \rightarrow \mathbf{G}(\mathcal{X}).$$

which corresponds under the isomorphisms of Corollary 3.4 to a map

$$\mathbf{G}(\mathcal{Y}_{\text{cl}}) \rightarrow \mathbf{G}(\mathcal{X}_{\text{cl}})$$

which can be thought of as a “virtual pullback” or virtual Gysin map along the morphism of classical stacks $f_{\text{cl}} : \mathcal{X}_{\text{cl}} \rightarrow \mathcal{Y}_{\text{cl}}$. It agrees with the virtual pullbacks of [Qu] (as will follow from the next subsection).

6.2. Fundamental classes via deformation to the normal stack. In this subsection we give an alternative construction of the maps $f^* : \mathbf{G}(\mathcal{Y}) \rightarrow \mathbf{G}(\mathcal{X})$ in the case when $f : \mathcal{X} \rightarrow \mathcal{Y}$ is quasi-smooth. This description was implicitly used in the proof of the Grothendieck–Riemann–Roch formula proven in [Kh5] (see Subject. 6.3 below).

Recall that for any quasi-smooth morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ there is a canonical “deformation to the normal stack” which sits in a commutative diagram

$$\begin{array}{ccccc} \mathcal{X} & \longrightarrow & \mathcal{X} \times \mathbf{A}^1 & \longleftarrow & \mathcal{X} \times \mathbf{G}_m \\ \downarrow 0 & & \downarrow & & \downarrow f \times \text{id} \\ N_{\mathcal{X}/\mathcal{Y}} & \xrightarrow{\hat{i}} & D_{\mathcal{X}/\mathcal{Y}} & \xleftarrow{\hat{j}} & \mathcal{Y} \times \mathbf{G}_m \\ \downarrow v & & \downarrow u & & \parallel \\ \mathcal{Y} & \xrightarrow{i} & \mathcal{Y} \times \mathbf{A}^1 & \xleftarrow{j} & \mathcal{Y} \times \mathbf{G}_m \end{array} \quad (6.6)$$

This is a family of quasi-smooth morphisms $\mathcal{X} \times \mathbf{A}^1 \rightarrow D_{\mathcal{X}/\mathcal{Y}}$ parametrized by the affine line whose generic fibre is the morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$ and whose special fibre is the zero section $0 : \mathcal{X} \rightarrow N_{\mathcal{X}/\mathcal{Y}}$ of the normal stack. The latter is by definition the vector bundle stack $\mathbf{V}(\mathcal{L}_{\mathcal{X}/\mathcal{Y}}[-1])$ associated to the (-1) -shifted cotangent complex. When f is a closed immersion, $N_{\mathcal{X}/\mathcal{Y}}$ is just the normal bundle and $D_{\mathcal{X}/\mathcal{Y}}$ is the usual deformation to the normal bundle. Note that the left-hand arrow v is the composite of the projection $\pi : N_{\mathcal{X}/\mathcal{Y}} \rightarrow \mathcal{X}$ with $f : \mathcal{X} \rightarrow \mathcal{Y}$. See [Kh5, Thm. 1.3].

Using this deformation we can construct a specialization map in G-theory in parallel to the case of Borel–Moore homology in [Kh5]. Consider the localization triangle (Theorem 3.9) associated to the lower row of (6.6):

$$\mathbf{G}(\mathcal{X} \times \mathbf{G}_m)[-1] \xrightarrow{\partial} \mathbf{G}(\mathcal{X}) \xrightarrow{i_*} \mathbf{G}(\mathcal{X} \times \mathbf{A}^1) \xrightarrow{j^*} \mathbf{G}(\mathcal{X} \times \mathbf{G}_m).$$

The boundary map ∂ admits a section

$$\gamma_b : \mathbf{G}(\mathcal{X}) \xrightarrow{q^*} \mathbf{G}(\mathcal{X} \times \mathbf{G}_m) \xrightarrow{\cap b} \mathbf{G}(\mathcal{X} \times \mathbf{G}_m)[-1]$$

where $q : \mathcal{X} \times \mathbf{G}_m \rightarrow \mathcal{X}$ is the projection and $b \in \mathbf{K}(\mathcal{X} \times \mathbf{G}_m)[-1]$ is the Bott class (inverse image of $b \in \mathbf{K}(\mathbf{Z}[T^{\pm 1}])[-1]$).

Consider also the localization triangle associated to the middle row of (6.6). These fit into a commutative diagram

$$\begin{array}{ccccccc} \mathrm{G}(\mathcal{Y} \times \mathbf{G}_m)[-1] & \xrightarrow{\partial} & \mathrm{G}(\mathcal{Y}) & \xrightarrow{i_*} & \mathrm{G}(\mathcal{Y} \times \mathbf{A}^1) & \xrightarrow{j^*} & \mathrm{G}(\mathcal{Y} \times \mathbf{G}_m) \\ \parallel & & \downarrow v^* & & \downarrow u^* & & \parallel \\ \mathrm{G}(\mathcal{Y} \times \mathbf{G}_m)[-1] & \xrightarrow{\hat{\partial}} & \mathrm{G}(N_{\mathcal{X}/\mathcal{Y}}) & \xrightarrow{\hat{i}_*} & \mathrm{G}(D_{\mathcal{X}/\mathcal{Y}}) & \xrightarrow{\hat{j}^*} & \mathrm{G}(\mathcal{Y} \times \mathbf{G}_m) \end{array}$$

where ∂ and $\hat{\partial}$ are the respective boundary maps. The right-hand square commutes by functoriality of inverse image, the middle square commutes by the base change formula, and the left-hand square commutes as a consequence.

Definition 6.7. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-smooth morphism of noetherian derived Artin stacks. The *specialization map* is the composite

$$\mathrm{sp}_{\mathcal{X}/\mathcal{Y}} : \mathrm{G}(\mathcal{Y}) \xrightarrow{\gamma_b} \mathrm{G}(\mathcal{Y} \times \mathbf{G}_m)[-1] \xrightarrow{\hat{\partial}} \mathrm{G}(N_{\mathcal{X}/\mathcal{Y}}),$$

where the notation is as above.

Since γ_b is a section of ∂ , commutativity of the diagram above immediately yields a canonical identification

$$\mathrm{sp}_{\mathcal{X}/\mathcal{Y}} = v^* = \pi^* \circ f^*,$$

where $\pi : N_{\mathcal{X}/\mathcal{Y}} \rightarrow \mathcal{X}$ is the projection. As long as we assume \mathcal{X} has affine stabilizers (e.g. is a derived Deligne–Mumford stack), we know that the map π^* is invertible by homotopy invariance (Corollary 5.6). Thus we find that the inverse image map f^* can be described in terms of the specialization map:

Proposition 6.8. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-smooth morphism of noetherian derived Artin stacks with affine stabilizers. Then $f^* : \mathrm{G}(\mathcal{Y}) \rightarrow \mathrm{G}(\mathcal{X})$ is canonically homotopic to the composite*

$$\mathrm{G}(\mathcal{Y}) \xrightarrow{\mathrm{sp}_{\mathcal{X}/\mathcal{Y}}} \mathrm{G}(N_{\mathcal{X}/\mathcal{Y}}) \xrightarrow{(\pi^*)^{-1}} \mathrm{G}(\mathcal{X}).$$

Remark 6.9. Throughout the above discussion we can replace G-theory by rational étale G-theory. In that case we do not need to assume affineness of stabilizers, since homotopy invariance for vector bundle stacks always holds in $\mathrm{G}^{\acute{e}t}$ (Corollary 5.6).

6.3. Grothendieck–Riemann–Roch formulas. Let k be a field and \mathcal{X} a derived Artin stack of finite type over k . Via stable motivic homotopy theory (see [Ri], [Kh5]) one can construct canonical isomorphisms

$$\tau_{\mathcal{X}} : \mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}} \rightarrow \bigoplus_{n \in \mathbf{Z}} \mathbf{R}\Gamma^{\mathrm{BM}}(\mathcal{X}, \mathbf{Q}(n))[-2n],$$

where on the right-hand side are spectra whose homotopy groups are the motivic Borel–Moore homology groups. Combining with the étale localization map $\mathrm{G}(\mathcal{X})_{\mathbf{Q}} \rightarrow \mathrm{G}^{\acute{e}t}(\mathcal{X})_{\mathbf{Q}}$, we get a map

$$\tau_{\mathcal{X}} : \mathrm{G}(\mathcal{X})_{\mathbf{Q}} \rightarrow \bigoplus_{n \in \mathbf{Z}} \mathbf{R}\Gamma^{\mathrm{BM}}(\mathcal{X}, \mathbf{Q}(n))[-2n]$$

which on π_0 gives a map

$$\tau_{\mathcal{X}} : G_0(\mathcal{X})_{\mathbf{Q}} \rightarrow H_{2n}^{\text{BM}}(\mathcal{X}, \mathbf{Q}(n)) \simeq A_*(\mathcal{X}_{\text{cl}})_{\mathbf{Q}}$$

where, if \mathcal{X} has affine stabilizers, the target is now identified with the Chow groups of the classical truncation \mathcal{X}_{cl} . This is a derived and stacky extension of the Baum–Fulton–MacPherson transformation [BFM] and we have the following analogue of the main result of *op. cit.*

Theorem 6.10.

- (i) *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper morphism between derived Artin stacks over k . Then there is a commutative square*

$$\begin{array}{ccc} G_0^{\text{ét}}(\mathcal{X})_{\mathbf{Q}} & \xrightarrow{f_*} & G_0^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}} \\ \downarrow \tau_{\mathcal{X}} & & \downarrow \tau_{\mathcal{Y}} \\ H_{2n}^{\text{BM}}(\mathcal{X}, \mathbf{Q}(n)) & \xrightarrow{f_*} & H_{2n}^{\text{BM}}(\mathcal{Y}, \mathbf{Q}(n)). \end{array}$$

- (ii) *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper representable morphism between derived Artin stacks over k . Then there is a commutative square*

$$\begin{array}{ccc} G_0(\mathcal{X})_{\mathbf{Q}} & \xrightarrow{f_*} & G_0(\mathcal{Y})_{\mathbf{Q}} \\ \downarrow \tau_{\mathcal{X}} & & \downarrow \tau_{\mathcal{Y}} \\ H_{2n}^{\text{BM}}(\mathcal{X}, \mathbf{Q}(n)) & \xrightarrow{f_*} & H_{2n}^{\text{BM}}(\mathcal{Y}, \mathbf{Q}(n)). \end{array}$$

Proof. For the first claim, one may identify $G_0^{\text{ét}}(\mathcal{X})$ with $\mathbf{R}\Gamma^{\text{BM}}(\mathcal{X}, \text{KGL})$ in a way that is natural with respect to proper direct images (see [Kh5, Ex. 2.13]). Then the claim follows immediately from the construction of proper direct images in Borel–Moore homology theories given in [Kh5]. The second claim follows by composing with the first with the square

$$\begin{array}{ccc} G_0(\mathcal{X})_{\mathbf{Q}} & \xrightarrow{f_*} & G_0(\mathcal{Y})_{\mathbf{Q}} \\ \downarrow \tau_{\mathcal{X}} & & \downarrow \tau_{\mathcal{Y}} \\ G_0^{\text{ét}}(\mathcal{X})_{\mathbf{Q}} & \xrightarrow{f_*} & G_0^{\text{ét}}(\mathcal{Y})_{\mathbf{Q}} \end{array}$$

which commutes by construction when f is representable (see Subsect. 5.4). \square

Remark 6.11. The representability hypothesis in claim (ii) is necessary, see Warning 5.10. However, following [To1], it is possible to prove a variant where the lower horizontal map is replaced by the direct image of the induced morphism of inertia stacks. See [KPRY].

In [Kh5] a fundamental class $[\mathcal{X}]$ is constructed in motivic Borel–Moore homology, when \mathcal{X} is quasi-smooth, and the following comparison with the G-theoretic fundamental class $[\mathcal{X}]_{\text{G}} = [\mathcal{O}_{\mathcal{X}}] \in G(\mathcal{X})$ (conjectured in [To3, Question 4.7]) is proven.

Theorem 6.12. *For every quasi-smooth derived Artin stack \mathcal{X} over k , there is an equality*

$$[\mathcal{X}] = \mathrm{Td}_{\mathcal{X}}^{-1} \cap \tau_{\mathcal{X}}[\mathcal{X}]_{\mathrm{G}},$$

where $\mathrm{Td}_{\mathcal{X}}$ is the Todd class of the cotangent complex $\mathcal{L}_{\mathcal{X}}$.

Remark 6.13. Via the identification mentioned above we may view the fundamental class $[\mathcal{X}]$ as a class $[\mathcal{X}_{\mathrm{cl}}]_{\mathrm{vir}} \in \mathbf{A}_*(\mathcal{X}_{\mathrm{cl}})_{\mathbf{Q}}$. For \mathcal{X} Deligne–Mumford, this coincides with the virtual fundamental class of Behrend–Fantechi [BF] (with respect to obstruction theory on $\mathcal{X}_{\mathrm{cl}}$ induced by the cotangent complex of \mathcal{X}) as explained in [Kh5, §3.3]. We can then read Theorem 6.12 as a comparison with the G-theoretic virtual fundamental class (Remark 6.3) via the formula

$$[\mathcal{X}_{\mathrm{cl}}]_{\mathrm{vir}} = (\mathrm{Td}_{\mathcal{X}_{\mathrm{cl}}}^{\mathrm{vir}})^{-1} \cap \tau_{\mathcal{X}_{\mathrm{cl}}}([\mathcal{X}_{\mathrm{cl}}]_{\mathrm{G}}^{\mathrm{vir}})$$

in $\mathbf{A}_*(\mathcal{X}_{\mathrm{cl}})_{\mathbf{Q}}$, where $\mathrm{Td}_{\mathcal{X}_{\mathrm{cl}}}^{\mathrm{vir}}$ is the Todd class of $\mathcal{L}_{\mathcal{X}}|_{\mathcal{X}_{\mathrm{cl}}}$. This recovers the virtual Grothendieck–Riemann–Roch formulas of [FG, CFK, LS], all proven in the case where \mathcal{X} is a derived scheme admitting an embedding into a smooth ambient scheme, as well as the extension to the case of quotient stacks in [RS].

6.4. K-theoretic fundamental classes. The following construction is a K-theory analogue of the “cohomological fundamental class” studied in [Kh5, §3.4].

Construction 6.14. Let \mathcal{X} be a derived Artin stack. For every quasi-smooth proper representable morphism $f : \mathcal{Z} \rightarrow \mathcal{X}$, there is a canonical class

$$[\mathcal{Z}/\mathcal{X}]^{\mathrm{K}} := f_*(1) = [f_*(\mathcal{O}_{\mathcal{Z}})] \in \mathbf{K}^{\mathrm{B}}(\mathcal{X}),$$

where $f_* : \mathbf{K}^{\mathrm{B}}(\mathcal{Z}) \rightarrow \mathbf{K}^{\mathrm{B}}(\mathcal{X})$ is the direct image map (which exists by Lemma 1.15) and $1 = [\mathcal{O}_{\mathcal{Z}}] \in \mathbf{K}^{\mathrm{B}}(\mathcal{Z})$ is the unit. If f is a closed immersion, then $[\mathcal{Z}/\mathcal{X}]^{\mathrm{K}}$ is supported on \mathcal{Z} , i.e., it lives in $\mathbf{K}^{\mathrm{B}}(\mathcal{X}$ on $\mathcal{Z})$.

When \mathcal{X} is nonsingular, this class is just the direct image of the G-theoretic fundamental class of \mathcal{Z} :

Remark 6.15. Let \mathcal{X} be a regular Artin stack. For any quasi-smooth proper representable morphism $f : \mathcal{Z} \rightarrow \mathcal{X}$, consider the commutative square

$$\begin{array}{ccc} \mathbf{K}^{\mathrm{B}}(\mathcal{Z}) & \xrightarrow{f_*} & \mathbf{K}^{\mathrm{B}}(\mathcal{X}) \\ \downarrow & & \downarrow \\ \mathbf{G}(\mathcal{Z}) & \xrightarrow{f_*} & \mathbf{G}(\mathcal{X}) \end{array}$$

where the vertical arrows are the Cartan maps (Theorem 3.5). Since \mathcal{X} is regular, the right-hand vertical arrow is invertible. Under this identification, it follows that the K-theoretic fundamental class $[\mathcal{Z}/\mathcal{X}]^{\mathrm{K}} \in \mathbf{K}^{\mathrm{B}}(\mathcal{X})$ can be identified with the direct image of the G-theoretic fundamental class $[\mathcal{Z}]_{\mathrm{G}} \in \mathbf{G}(\mathcal{Z})$ (Construction 6.2).

Proposition 6.16. *Suppose given a commutative square of derived Artin stacks*

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow p & & \downarrow q \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

where i and i' are quasi-smooth closed immersions. If the square is homotopy cartesian, then we have a canonical identification

$$q^*[\mathcal{Z}/\mathcal{X}] \simeq [\mathcal{Z}'/\mathcal{X}']$$

in $\mathbf{K}^{\mathbf{B}}(\mathcal{X}')$.

Proof. Evaluate the base change formula $q^*i_* \simeq i'_*p^*$ (Proposition 2.8) on the unit $1 = [\mathcal{O}_{\mathcal{Z}}] \in \mathbf{K}^{\mathbf{B}}(\mathcal{Z})$. \square

Proposition 6.17 (Non-transverse Bézout formula). *Let \mathcal{X} be a derived Artin stack. Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ and $g : \mathcal{Z} \rightarrow \mathcal{X}$ be quasi-smooth proper representable morphisms. Then we have*

$$[\mathcal{Y}/\mathcal{X}]^{\mathbf{K}} \cup [\mathcal{Z}/\mathcal{X}]^{\mathbf{K}} \simeq [\mathcal{Y} \times_{\mathcal{X}}^{\mathbf{R}} \mathcal{Z}/\mathcal{X}]^{\mathbf{K}}$$

in $\mathbf{K}^{\mathbf{B}}(\mathcal{X})$.

Proof. By the projection and base change formulas (Propositions 2.7 and 2.8), we have

$$f_*(1) \cup g_*(1) \simeq f_*f^*g_*(1) \simeq h_*(1)$$

where $h : \mathcal{Y} \times_{\mathcal{X}}^{\mathbf{R}} \mathcal{Z} \rightarrow \mathcal{X}$ is the projection. \square

To deal with non-proper intersections there is an *excess intersection formula* which expresses the failure of the base change formula (Proposition 6.16) in terms of the K-theoretic Euler class of the excess bundle, see [T4] and [Kh6]. For example:

Proposition 6.18 (Self-intersection formula). *Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a quasi-smooth closed immersion. Then there is a canonical identification*

$$i^*i_*(-) \simeq e(\mathcal{N}_{\mathcal{Z}/\mathcal{X}}) \cup (-)$$

of maps $\mathbf{K}^{\mathbf{B}}(\mathcal{Z}) \rightarrow \mathbf{K}^{\mathbf{B}}(\mathcal{X})$. In particular,

$$i^*[\mathcal{Z}/\mathcal{X}] \simeq e(\mathcal{N}_{\mathcal{Z}/\mathcal{X}})$$

in $\mathbf{K}^{\mathbf{B}}(\mathcal{X})$ and

$$[\mathcal{Z}/\mathcal{X}] \cup [\mathcal{Z}/\mathcal{X}] \simeq [\mathcal{Z} \times_{\mathcal{X}}^{\mathbf{R}} \mathcal{Z}/\mathcal{X}] \simeq i_*(e(\mathcal{N}_{\mathcal{Z}/\mathcal{X}}))$$

in $\mathbf{K}^{\mathbf{B}}(\mathcal{X})$ (or $\mathbf{K}^{\mathbf{B}}(\mathcal{X}$ on $\mathcal{Z})$).

6.5. The γ -filtration and Chow cohomology of singular schemes.

This subsection is a digression that attempts to justify my interest in these “K-theoretic fundamental classes” (Subsect. 6.4), which do not seem to have received much attention in the literature so far.

6.5.1. *The γ -filtration and Chow cohomology of nonsingular schemes.* Let X be a scheme of finite type over a field k and consider the group $Z_*(X)$ of algebraic cycles on X . Recall that there is a map

$$Z_*(X) \rightarrow G_0(X), \quad [Z] \mapsto [\mathcal{O}_Z].$$

This map does not respect rational equivalence but does so if we pass from $G_0(X)$ to the graded pieces of the coniveau or γ -filtration (both agree with rational coefficients); there are canonical surjections

$$A_i(X) \rightarrow \mathrm{Gr}^{d-i}(G_0(X)), \quad [Z] \mapsto [\mathcal{O}_Z]$$

where $A_*(X)$ denotes the quotient of $Z_*(X)$ modulo rational equivalence.

When X is nonsingular, the Grothendieck–Riemann–Roch theorem implies that the kernel of this map is torsion. Moreover the γ -filtration is compatible with the cup product on $K_0(X)$ and on the “Poincaré dual” theories there is an isomorphism of graded rings

$$A^*(X)_{\mathbf{Q}} \rightarrow \mathrm{Gr}^*(K_0(X))_{\mathbf{Q}}. \quad (6.19)$$

Here $A^*(X) = A_{d-*}(X)$ by definition and $K_0(X) \simeq G_0(X)$ by Theorem 3.5. See [SGA6, Exp. 0, Appendix and Exp. 14, §4] and [Fu2, Ex. 15.3.6].

6.5.2. *Chow cohomology of singular schemes.* For singular X the right-hand side of (6.19) still makes sense but it is less clear how to define “Chow cohomology” rings $A^*(X)$ such that (6.19) holds.⁶ There is a reasonable definition of top-degree Chow cohomology $A^d(X)$, where X is a singular d -dimensional variety, due to Levine and Weibel, see e.g. [LW, Sr, GK]. Here I want to briefly explain how the theory of K-theoretic fundamental classes suggests a construction of a “derived” Chow cohomology theory.

Let $Z_{\mathrm{der}}^n(X)$ denote the free abelian group on quasi-smooth projective morphisms $f : Z \rightarrow X$ of relative virtual dimension $-n$. Denote the class of $f : Z \rightarrow X$ by $[Z/X] \in Z_{\mathrm{der}}^n(X)$. There is an obvious product

$$[Z/X] \cup [Z'/X] = [Z \times_X^{\mathbf{R}} Z'/X],$$

as well as a map

$$Z_{\mathrm{der}}^*(X) \rightarrow K_0(X), \quad [Z/X] \mapsto [Z/X]^{\mathbf{K}}$$

⁶Fulton’s operational Chow groups [Fu2, Chap. 17] are commonly used as a substitute for a “genuine” Chow cohomology theory, but they satisfy neither this property nor many others (for example they do not even map to singular cohomology, see [Tot2]). Voevodsky’s motivic cohomology groups [Vo] are almost what one is looking for except that rationally they compare to the *homotopy invariant* version of K-theory. On singular schemes, motivic cohomology can be computed (as a presheaf of complexes) by the following procedure: restrict to smooth k -schemes, take the left Kan extension to k -schemes of finite type, and finally take the \mathbf{A}^1 -homotopy localization of the result (see [Kh7, Prop. 5]). Stopping before the last step gives a non-homotopy invariant version of motivic cohomology. This is a variant of an old construction of Fulton from [Fu1, §3.1]. A similar procedure works for homotopy invariant algebraic K-theory (see e.g. [CiKh, Props. 5.2.2 and 5.3.7]) and in that case stopping before the last step does recover algebraic K-theory (see [Fu1, §3.2], [EHKSY, App. A]). The derived Chow ring constructed below should hopefully compute these non-homotopy invariant motivic cohomology groups for nice enough schemes.

where $[Z/X]^K = [f_*(\mathcal{O}_Z)]$ is the K-theoretic fundamental class (Subsect. 6.4). There should be some natural quotient $Z_{\text{der}}^*(X) \twoheadrightarrow A_{\text{der}}^*(X)$ and an isomorphism

$$A_{\text{der}}^*(X)_{\mathbf{Q}} \rightarrow \text{Gr}^*(K_0(X))_{\mathbf{Q}}. \quad (6.20)$$

The following is a sort of sanity check for this hypothesis.

Theorem 6.21. *For every qcqs algebraic space X and every $[Z/X] \in Z_{\text{der}}^n(X)$, we have*

$$[Z/X]^K \in \text{Gr}^n(K_0(X))_{\mathbf{Q}}.$$

More generally, the map $f_* : K_0(Z) \rightarrow K_0(X)$ sends $\text{Fil}^k K_0(Z)_{\mathbf{Q}}$ to $\text{Fil}^{k+n} K_0(X)_{\mathbf{Q}}$ for every k .

Proof. By embedding Z into a projective bundle over X and using the projective bundle formula one reduces to the case of a quasi-smooth closed immersion. That case is proven in [Kh3, Thm. 2] using derived blow-ups [KhRy] and the excess intersection formula [Kh6] to further reduce to the case of virtual Cartier divisors. \square

In particular, there is an induced Gysin map $f_*^\gamma : \text{Gr}^* K_0(Z)_{\mathbf{Q}} \rightarrow \text{Gr}^* K_0(X)_{\mathbf{Q}}$. One has moreover the following version of the Grothendieck–Riemann–Roch theorem, a direct generalization of the original formulation in [SGA6, Exp. VIII, Thm. 3.6]:

Theorem 6.22. *Let X be a qcqs algebraic space. For every quasi-smooth projective morphism $f : Z \rightarrow X$ of relative virtual dimension n , there is a commutative square*

$$\begin{array}{ccc} K_0(Z) & \xrightarrow{f_*} & K_0(X) \\ \downarrow \text{ch} & & \downarrow \text{ch} \\ \text{Gr}^* K_0(Z)_{\mathbf{Q}} & \xrightarrow{f_*^\gamma(-\cdot \text{Td}(\mathcal{L}_{Z/X}^\vee))} & \text{Gr}^* K_0(X)_{\mathbf{Q}} \end{array}$$

where $\text{Td}(\mathcal{L}_{Z/X}^\vee)$ is the Todd class of the relative tangent complex.

Proof. Again, one reduces to the case of projective bundles, see [SGA6, Exp. VIII, §5], and the case of quasi-smooth closed immersions, see [Kh3, Thm. 3]. \square

The theory of fundamental classes in motivic cohomology [Kh5] produces a ring homomorphism

$$Z_{\text{der}}^*(X) \rightarrow H^{2*}(X, \mathbf{Z}_{\text{mot}}(*))$$

which should descend to $A_{\text{der}}^*(X)$. When X is smooth the resulting map

$$A_{\text{der}}^*(X) \rightarrow H^{2*}(X, \mathbf{Z}_{\text{mot}}(*)) \simeq A^*(X) \quad (6.23)$$

should be an isomorphism. Note that if the base field has resolution of singularities then every cycle in $A^n(X)$ can be represented by the fundamental class of a projective lci map, so there is at least some quotient of $Z_{\text{der}}^*(X)$ for which (6.23) is an isomorphism. Whenever this is true, it means the

intersection product on $A^*(X)$ can be represented geometrically by derived fibred products.

Admittedly A_{der}^* looks more like a cobordism theory than a cycle theory. In fact two completely different recent approaches to algebraic cobordism both yield constructions closely resembling $A_{\text{der}}^*(X)$. In [EHKSY] the authors considered a variant where the generators are required to be *finite* instead of projective. That theory gives a model for cobordism of *smooth* schemes and unfortunately requires a Nisnevich localization procedure which renders the end result fairly intractible. On the other hand Annala has recently constructed good theories of algebraic cobordism and Chow cohomology of singular schemes in characteristic zero [An1]. After further simplification and partial extension to general bases in [AnYo, An2], the generators are similar to those of A_{der}^* and, according to a private communication from Annala, there is a simple set of relations one can impose on his “precobordism” such that (6.20) holds (under very mild hypotheses on X). Comparisons with the Levine–Weibel Chow groups [LW] have yet to be investigated.

REFERENCES

- [AGH] B. Antieau, D. Gepner, J. Heller, *K-theoretic obstructions to bounded t-structures*. Invent. Math. **216** (2019), no. 1, 241–300.
- [AHR1] J. Alper, J. Hall, D. Rydh, *A Luna étale slice theorem for algebraic stacks*, Ann. Math. (2) **191**, no. 3, 675–738 (2020).
- [AHR2] J. Alper, J. Hall, D. Rydh, *The étale local structure of algebraic stacks*, arXiv:1912.06162 (2019).
- [AHRH] J. Alper, J. Hall, D. Halpern-Leistner, D. Rydh, *Artin algebraization for pairs with applications to the local structure of stacks and Ferrand pushouts*. In preparation.
- [An1] T. Annala, *Bivariant derived algebraic cobordism*. J. Algebr. Geom., to appear. arXiv:1807.04989 (2018).
- [An2] T. Annala, *Precobordism and cobordism*. arXiv:2006.11723 (2020).
- [AnYo] T. Annala, S. Yokura, *Bivariant algebraic cobordism with bundles*. arXiv:1911.12484 (2019).
- [AOV] D. Abramovich, M. Olsson, A. Vistoli, *Tame stacks in positive characteristic*. Ann. Inst. Fourier **58**, no. 4, 1057–1091 (2008); corrigendum ibid. **64**, no. 3, 945–946 (2014).
- [B] H. Bass, *Algebraic K-theory*. Math. Lect. Note Series (1968).
- [BFN] D. Ben-Zvi, J. Francis, D. Nadler, *Integral transforms and Drinfeld centers in derived algebraic geometry*. J. Am. Math. Soc. **23** (2010), no. 4, 909–966.
- [Ba1] C. Barwick, *On exact ∞ -categories and the theorem of the heart*. Compos. Math. **151** (2015), no. 11, 2160–2186.
- [Ba2] C. Barwick, *On the algebraic K-theory of higher categories*. J. Topol. **9**, no. 1, 245–347 (2016).
- [BF] K. Behrend, B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128** (1997), no. 1, 45–88.
- [BFM] P. Baum, W. Fulton, R. MacPherson, *Riemann–Roch for singular varieties*. Publ. Math. IHÉS **45**, 101–145 (1975).
- [BGT] A. J. Blumberg, D. Gepner, G. Tabuada, *A universal characterization of higher algebraic K-theory*. Geom. Topol. **17** (2013), no. 2, 733–838.
- [BKRS] T. Bachmann, A. A. Khan, C. Ravi, V. Sosnilo, *Categorical Milnor squares and algebraic K-theory of stacks*. arXiv:2011.04355 (2020).
- [BS] B. Bhatt, P. Scholze, *Projectivity of the Witt vector affine Grassmannian*. Invent. Math. **209** (2017), no. 2, 329–423.

- [BVdB] A. Bondal, M. van den Bergh, *Mosc. Math. J.* **3**, no. 1, 1–36 (2003).
- [CD] D.-C. Cisinski, F. Déglise, *Triangulated categories of mixed motives*. Springer Monogr. Math., to appear. [arXiv:0912.2110](https://arxiv.org/abs/0912.2110) (2012).
- [CFK] I. Ciocan-Fontanine, M. Kapranov, *Virtual fundamental classes via dg-manifolds*. *Geom. Topol.* **13** (2009), no. 3, 1779–1804.
- [Ci] D.-C. Cisinski, *Descente par éclatements en K -théorie invariante par homotopie*. *Ann. of Math.* **177** (2013), no. 2, 425–448.
- [CiKh] D.-C. Cisinski, A. A. Khan, \mathbf{A}^1 -homotopy invariance in spectral algebraic geometry. [arXiv:1705.03340](https://arxiv.org/abs/1705.03340) (2020).
- [CMNN] D. Clausen, A. Mathew, N. Naumann, J. Noel, *Descent in algebraic K -theory and a conjecture of Ausoni–Rognes*. *J. Eur. Math. Soc. (JEMS)* **22** (2020), no. 4, 1149–1200.
- [DG] V. Drinfeld, D. Gaitsgory, *On some finiteness questions for algebraic stacks*. *Geom. Funct. Anal.* **23**, no. 1, 149–294 (2013).
- [EHKSY] E. Elmanto, M. Hoyois, A. A. Khan, V. Sosnilo, M. Yakerson, *Modules over algebraic cobordism*. *Forum Math. Pi*, to appear. [arXiv:1908.02162](https://arxiv.org/abs/1908.02162).
- [EGA] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique*, Publ. Math. IHÉS **4** (Chapter 0, 1–7, and I, 1–10), **8** (II, 1–8), **11** (Chapter 0, 8–13, and III, 1–5), **17** (III, 6–7), **20** (Chapter 0, 14–23, and IV, 1), **24** (IV, 2–7), **28** (IV, 8–15), and **32** (IV, 16–21), 1960–1967.
- [Fa] G. Faltings, *Finiteness of coherent cohomology for proper fppf stacks*. *J. Algebr. Geom.* **12**, no. 2, 357–366 (2003).
- [FG] B. Fantechi, L. Göttsche, *Riemann–Roch theorems and elliptic genus for virtually smooth schemes*. *Geom. Topol.* **14**, no. 1, 83–115 (2010).
- [Fu1] W. Fulton, *Rational equivalence on singular varieties*, Publ. Math. IHÉS **45**, 147–167 (1975).
- [Fu2] W. Fulton, *Intersection theory*. Springer (1998).
- [GGN] D. Gepner, M. Groth, T. Nikolaus, *Universality of multiplicative infinite loop space machines*. *Algebr. Geom. Topol.* **15**, no. 6, 3107–3153 (2015).
- [GK] R. Gupta, A. Krishna, *K -theory and 0-cycles on schemes*. *J. Algebr. Geom.* **29**, no. 3, 547–601 (2020).
- [GR] D. Gaitsgory, N. Rozenblyum, *A study in derived algebraic geometry. Vol. I. Correspondences and duality*. *Math. Surv. Mono.* **221** (2017).
- [Gr] P. Gross, *Tensor generators on schemes and stacks*. *Algebr. Geom.* **4**, no. 4, 501–522 (2017).
- [HLP] D. Halpern-Leistner, A. Preygel, *Mapping stacks and categorical notions of properness*. [arXiv:1402.3204](https://arxiv.org/abs/1402.3204) (2019).
- [Ho1] M. Hoyois, *A quadratic refinement of the Grothendieck–Lefschetz–Verdier trace formula*, *Algebr. Geom. Topol.* **14** (2014), no. 6, 3603–3658.
- [Ho2] M. Hoyois, *The six operations in equivariant motivic homotopy theory*, *Adv. Math.* **305**, 197–279 (2017); corrigendum *ibid.* **333**, 1293–1296 (2018).
- [Ho3] M. Hoyois, *K -theory of dualizable categories (after A. Efimov)*. <http://www.mathematik.ur.de/hoyois/papers/efimov.pdf> (2018).
- [Ho4] M. Hoyois, *Cdh descent in equivariant homotopy K -theory*. *Doc. Math.* **25**, 457–482 (2020).
- [HoKr] M. Hoyois, A. Krishna, *Vanishing theorems for the negative K -theory of stacks*. *Ann. K-Theory* **4**, no. 3, 439–472 (2019).
- [HR1] J. Hall, D. Rydh, *Algebraic groups and compact generation of their derived categories of representations*. *Indiana Univ. Math. J.* **64**, no. 6, 1903–1923 (2015).
- [HR2] J. Hall, D. Rydh, *Perfect complexes on algebraic stacks*. *Compos. Math.* **153**, no. 11, 2318–2367 (2017).
- [Ji] F. Jin, *Algebraic G -theory in motivic homotopy categories*. [arXiv:1806.03927](https://arxiv.org/abs/1806.03927) (2018).
- [Jo] R. Joshua, *Riemann–Roch for algebraic stacks. I*. *Compositio Math.* **136** (2003), no. 2, 117–169.

- [Kh1] A. A. Khan, *The Morel–Voevodsky localization theorem in spectral algebraic geometry*, *Geom. Topol.* **23** (2019), no. 7, 3647–3685. [arXiv:1610.06871](https://arxiv.org/abs/1610.06871).
- [Kh2] A. A. Khan, *Pro-systems of K-theory spectra*. Lecture notes, available at <https://www.preschema.com/lecture-notes/kdescent/lect8.pdf> (2017).
- [Kh3] A. A. Khan, *The Grothendieck–Riemann–Roch theorem*. Lecture notes, available at <https://www.preschema.com/teaching/grr-ss18/lect10.pdf> (2018).
- [Kh4] A. A. Khan, *Algebraic K-theory of quasi-smooth blow-ups and cdh descent*. *Ann. H. Lebesgue*, to appear. [arXiv:1810.12858](https://arxiv.org/abs/1810.12858).
- [Kh5] A. A. Khan, *Virtual fundamental classes for derived stacks I*. [arXiv:1909.01332](https://arxiv.org/abs/1909.01332) (2019).
- [Kh6] A. A. Khan, *Virtual excess intersection theory*. [arXiv:1909.13829](https://arxiv.org/abs/1909.13829) (2019).
- [Kh7] A. A. Khan, *The cdh-local motivic homotopy category (after Cisinski)*. <https://www.preschema.com/papers/shcdh.pdf> (2019).
- [KhRa] A. A. Khan, C. Ravi, *Generalized cohomology theories for algebraic stacks*. [arXiv:2106.15001](https://arxiv.org/abs/2106.15001) (2021).
- [KhRy] A. A. Khan, D. Rydh, *Virtual Cartier divisors and blow-ups*. [arXiv:1802.05702](https://arxiv.org/abs/1802.05702) (2018).
- [KPRY] A. A. Khan, M. Porta, C. Ravi, T. Y. Yu. In preparation.
- [Ko] M. Kontsevich, *Enumeration of rational curves via torus actions*, in: *The moduli space of curves (Texel Island, 1994)*, 335–368, *Progr. Math.*, **129** (1995).
- [KrRa] A. Krishna, C. Ravi, *Algebraic K-theory of quotient stacks*. *Ann. K-Theory* **3**, no. 2, 207–233 (2018).
- [KST] M. Kerz, F. Strunk, G. Tamme, *Algebraic K-theory and descent for blow-ups*. *Invent. Math.* **211**, no. 2, 523–577 (2018).
- [Lee] Y.-P. Lee, *Quantum K-theory, I: Foundations*. *Duke Math. J.* **121**, no. 3, 389–424 (2004).
- [LS] P. Lowrey, T. Schürig, *Grothendieck–Riemann–Roch for derived schemes*. [arXiv:1208.6325](https://arxiv.org/abs/1208.6325) (2012).
- [LT] M. Land, G. Tamme, *On the K-theory of pullbacks*, *Ann. Math. (2)* **190**, no. 3, 877–930 (2019).
- [Lu1] J. Lurie, *Higher topos theory*, *Ann. Math. Stud.* **170** (2009).
- [Lu2] J. Lurie, *Algebraic K-theory and manifold topology*, lecture notes (2014). <https://www.math.ias.edu/~lurie/281.html>.
- [Lu3] J. Lurie, *Higher algebra*, version of 2017-09-18. <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Lu4] J. Lurie, *Spectral algebraic geometry*, version of 2018-02-03. <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>.
- [LW] M. Levine, C. A. Weibel, *Zero cycles and complete intersections on singular varieties*, *J. Reine Angew. Math.* **359**, 106–120 (1985).
- [Ma] P. May, *E_∞ spaces, group completions, and permutative categories*, in: *New Developments in Topology*, 61–93, *Proc. Sympos. Algebr. Topology*, Oxford 1972 (1974).
- [Mi] J. W. Milnor, *Introduction to algebraic K-theory*. *Annals of Math. Studies*, no. 72 (1971).
- [MV] F. Morel, V. Voevodsky, *\mathbf{A}^1 -homotopy theory of schemes*. *Publ. Math. IHÉS* **90** (1999), 45–143.
- [Ol] M. C. Olsson, *On proper coverings of Artin stacks*, *Adv. Math.* **198**, no. 1, 93–106 (2005).
- [Q1] D. Quillen, *On the group completion of a simplicial monoid* (1971).
- [Q2] D. Quillen, *Higher algebraic K-theory: I*, in: *Cohomology of groups and algebraic K-theory*, 413–478, *Adv. Lect. Math. (ALM)* **12**, Int. Press (2010).
- [Qu] F. Qu, *Virtual pullbacks in K-theory*, *Ann. Inst. Fourier* **68**, no. 4, 1609–1641 (2018).
- [Ri] J. Riou, *Algebraic K-theory, \mathbf{A}^1 -homotopy and Riemann–Roch theorems*, *J. Topol.* **3** (2010), no. 2, 229–264.
- [RS] C. Ravi, B. Sreedhar, *Virtual equivariant Grothendieck–Riemann–Roch formula*. [arXiv:2009.09697](https://arxiv.org/abs/2009.09697) (2020).

- [Ry1] D. Rydh, *Submersions and effective descent of étale morphisms*. Bull. Soc. Math. France **138** (2010), no. 2, 181–230.
- [Ry2] D. Rydh, *Approximation of sheaves on algebraic stacks*. Int. Math. Res. Not. **2016** (2016), no. 3, 717–737.
- [SGA6] P. Berthelot, A. Grothendieck, L. Illusie, *Théorie des intersections and théorème de Riemann–Roch*. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6). Lecture Notes in Mathematics **225**, Springer (1971).
- [SP] The Stacks Project. <https://stacks.math.columbia.edu>.
- [Sa] V. Sadhu, *On the vanishing of relative negative K-theory*. J. Algebra Appl. **19** (2020), no. 8, Article ID 2050152, 14 p.
- [Spi] M. Spivakovsky, *A new proof of D. Popescu’s theorem on smoothing of ring homomorphisms*. J. Am. Math. Soc. **12**, no. 2, 381–444 (1999).
- [Spr] T. A. Springer, *Linear algebraic groups*. 2nd ed. Birkhäuser (1998).
- [Sr] V. Srinivas, *Zero cycles on singular varieties*, in: NATO ASI Ser., Ser. C, Math. Phys. Sci. **548**, 347–382 (2000).
- [Sw] R. G. Swan, *Excision in algebraic K-theory*. J. Pure Appl. Algebra **1**, 221–252 (1971).
- [T1] R. W. Thomason, *Algebraic K-theory of group scheme actions*, in: Algebraic topology and algebraic K-theory, Proc. Conf., Princeton, NJ (USA), Ann. Math. Stud. **113**, 539–563 (1987).
- [T2] R. W. Thomason, *Equivariant resolution, linearization, and Hilbert’s fourteenth problem over arbitrary base schemes*. Adv. Math. **65** (1987), 16–34.
- [T3] R. W. Thomason, *Les K-groupes d’un fibré projectif*, in: Algebraic K-theory and algebraic topology (Lake Louise, AB, 1991), 243–248, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., **407** (1993).
- [T4] R. W. Thomason, *Les K-groupes d’un schéma éclaté et une formule d’intersection excédentaire*, Invent. Math. **112**, 195–215 (1993).
- [Ta] G. Tamme, *Excision in algebraic K-theory revisited*. Compos. Math. **154** (2018), no. 9, 1801–1814.
- [To1] B. Toën, *Théorèmes de Riemann–Roch pour les champs de Deligne–Mumford*, K-Theory **18**, no. 1, 33–76 (1999).
- [To2] B. Toën, *Notes sur la G-théorie rationnelle des champs de Deligne–Mumford*. [arXiv:math/9912172](https://arxiv.org/abs/math/9912172) (1999).
- [To3] B. Toën, *Higher and derived stacks: a global overview*, Proc. Symp. Pure Math. **80**, 435–487 (2009). [arXiv:math/0604504](https://arxiv.org/abs/math/0604504).
- [To4] B. Toën, *Derived Azumaya algebras and generators for twisted derived categories*. Invent. Math. **189** (2012), no. 3, 581–652.
- [To5] B. Toën, *Derived algebraic geometry*, EMS Surv. Math. Sci. **1** (2014), 153–240.
- [Tot1] B. Totaro, *The resolution property for schemes and stacks*. J. Reine Angew. Math. **577**, 1–22 (2004).
- [Tot2] B. Totaro, *Chow groups, Chow cohomology, and linear varieties*. Forum Math. Sigma **2**, e17 (2014).
- [TV] B. Toën, M. Vaquié, *Moduli of objects in dg-categories*. Ann. Sci. Éc. Norm. Supér. **40** (2007), no. 3, 387–444.
- [TT] R. W. Thomason, T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, in: The Grothendieck Festschrift, Vol. III, 247–435, Progr. Math. **88**, Birkhäuser (1990).
- [Vo] V. Voevodsky, *Triangulated categories of mixed motives over a field*, Ann. Math. Stud. **143**, 188–238 (2000).
- [We1] C. A. Weibel, *K-theory and analytic isomorphisms*. Invent. Math. **61**, 177–197 (1980).
- [We2] C. A. Weibel, *Homotopy algebraic K-theory*, in: Algebraic K-theory and algebraic number theory, Proc. Semin., Honolulu/Hawaii 1987, Contemp. Math. **83**, 461–488 (1989).

Taipei
10617 Taiwan