

Exercise sheet 1

Prove all the propositions and/or theorems below.

Adjoint Functors.

Proposition 1. *Let $F : \mathcal{C} \rightleftarrows \mathcal{D} : G$ be an adjunction. Assume that the right adjoint G is fully faithful. If ever \mathcal{C} is complete (cocomplete), so is \mathcal{D} .*

Given a functor between small categories $u : A \rightarrow B$, we will write the functor of composition with u

$$u^* : \widehat{B} \rightarrow \widehat{A}.$$

It has a left adjoint

$$u_! : \widehat{A} \rightarrow \widehat{B}$$

obtained as the extension by colimits of the composition of u with the Yoneda embedding of B . In other words, the functor u is the unique functor which commutes with small colimits such that $u_!(h_a) = h_{u(a)}$ for any object a of A . One checks that the functor u^* also preserves small colimits, and thus has a right adjoint

$$u_* : \widehat{A} \rightarrow \widehat{B}$$

Proposition 2. *The following three conditions are equivalent.*

- (i) *The functor u is fully faithful.*
- (ii) *The functor $u_!$ is fully faithful.*
- (iii) *The functor u_* is fully faithful.*

Skeletons. We may apply the preceding construction in the following case. For any integer $n \geq 0$, let $\Delta_{\leq n}$ be the full subcategory whose objects are the sets of the form $[k]$ for $k \leq n$, and let $i_n : \Delta_{\leq n} \rightarrow \Delta$ be the inclusion functor. We thus have a sequence of adjoints $(i_n)_!$, i_n^* and $(i_n)_*$, and thus a pair of adjoint. One defines a pair of adjoint functors by setting, for any simplicial set:

$$sk_n(X) = (i_n)_!(i_n^*(X)) \quad \text{and} \quad cosk_n(X) = (i_n)_*(i_n^*(X)).$$

Note that, *a priori*, the simplicial set $sk_n(X)$ has to be distinguished from the skeleton $Sk_n(X)$ (which is defined as a subobject of X).

Proposition 3. *If, for a given integer $n \geq 0$, a morphism of simplicial sets $X \rightarrow Y$ induces injections $X_k \rightarrow Y_k$ for all $k \leq n$, and if $Sk_n(X) = X$, then it is a monomorphism.*

The preceding proposition is useful to prove:

Proposition 4. *The co-unit morphism $sk_n(X) \rightarrow X$ induces a canonical isomorphism*

$$sk_n(X) \simeq Sk_n(X).$$

Groupoids. Let Gpd denote the full subcategory of Cat spanned by small groupoids, and let $\iota : Gpd \hookrightarrow Cat$ denote the inclusion.

Proposition 5. *The functor ι admits a left adjoint π_1 and a right adjoint k .*

The right adjoint sends any category C to $k(C)$, its maximal subgroupoid. This is defined as the subcategory of C with the same objects and only the invertible morphisms.

The groupoid $\pi_1(C)$, sometimes called the *fundamental groupoid* of C , can be described as the localization of C at the set of all its morphisms (in the sense of Gabriel–Zisman).

Now consider the inclusion $i : Set \hookrightarrow Gpd$, where we view a set E as a groupoid $i(E)$ with $Ob(i(E)) = E$ and only identity morphisms.

Lemma 6. *The functor $i : Set \hookrightarrow Gpd$ admits a left adjoint π_0 .*

The set $\pi_0(K)$ is called the set of connected components of K .

Proposition 7. *For a small category C , the following conditions are equivalent.*

- (i) *The category C is a groupoid (i.e. all morphisms of C are invertible).*
- (ii) *The simplicial set $N(C)$ is an ∞ -groupoid.*
- (iii) *The simplicial set $N(C)$ is a Kan complex.*
- (iv) *There exists a presheaf X on the category of non-empty finite sets (with all maps as morphisms) whose restriction to $\mathbf{\Delta}$ is isomorphic to C .*