

## Exercise sheet 6

### 1. CATEGORIES OF CORRESPONDENCES

1.1. Recall that  $S(X)$  denotes the *twisted diagonal* of a simplicial set  $X$ , as in Sheet 5.

**Exercise 1.1.1.** Describe the  $\infty$ -category  $S(\Delta^n)$  for  $n \in \Delta$  (hint: it is the nerve of an ordinary category).

Below, it will be important to understand the shape of diagrams  $S(\Delta^n) \rightarrow \mathbf{C}$ .

1.2. Let  $\mathbf{C}$  be an  $\infty$ -category with two subcategories  $\mathbf{C}^{\text{coadm}}$  and  $\mathbf{C}_{\text{adm}}$ , with the following properties:

(i) Any isomorphism of  $\mathbf{C}$  is contained in  $\mathbf{C}^{\text{coadm}}$  and  $\mathbf{C}_{\text{adm}}$ . (Thus in particular each subcategory contains all objects of  $\mathbf{C}$ .)

(ii) For any morphism in  $\mathbf{C}^{\text{coadm}}$  (resp. in  $\mathbf{C}_{\text{adm}}$ ), the base change along a morphism in  $\mathbf{C}_{\text{adm}}$  (resp. in  $\mathbf{C}^{\text{coadm}}$ ) exists and belongs to  $\mathbf{C}^{\text{coadm}}$  (resp. to  $\mathbf{C}_{\text{adm}}$ ).

We will refer to a morphism of  $\mathbf{C}$  as *admissible* (resp. *co-admissible*) if it belongs to the subcategory  $\mathbf{C}^{\text{coadm}}$  (resp.  $\mathbf{C}_{\text{adm}}$ ). We will refer to a cartesian square in  $\mathbf{C}$  as *bi-admissible* if it is of the form

$$\begin{array}{ccc} c_{1,1} & \xrightarrow{f'} & c_{1,0} \\ \downarrow g' & & \downarrow g \\ c_{0,1} & \xrightarrow{f} & c_{0,0}, \end{array}$$

where  $f$  and  $f'$  are admissible, and  $g$  and  $g'$  are co-admissible.

1.3. Define a simplicial set

$$\text{Corr}(\mathbf{C}) \in \text{Set}_{\Delta}$$

whose  $n$ -simplices are functors  $\delta : S(\Delta^n)^{\text{op}} \rightarrow \mathbf{C}$ , satisfying the condition that each commutative square ( $0 \leq i \leq k \leq \ell \leq j \leq n$ )

$$\begin{array}{ccc} \mathbf{C}_{i,j} & \longrightarrow & \mathbf{C}_{k,j} \\ \downarrow & & \downarrow \\ \mathbf{C}_{i,\ell} & \longrightarrow & \mathbf{C}_{k,\ell}, \end{array}$$

appearing in the diagram  $\delta$ , is cartesian and bi-admissible.

**Exercise 1.3.1.** Describe the simplicial set  $\text{Corr}(\mathbf{C})$  in the case where  $\mathbf{C}$  is the nerve of an ordinary category.

**Exercise 1.3.2.** Describe the category  $\tau(\text{Corr}(\mathbf{C}))$ , where  $\mathbf{C}$  is an  $\infty$ -category.

**Exercise 1.3.3.** Identify the opposite simplicial set  $\text{Corr}(\mathbf{C})^{\text{op}}$ .

1.4. One would like to prove, among other things:

**Proposition 1.4.1.** *The simplicial set  $\text{Corr}(\mathbf{C})$  is an  $\infty$ -category.*

This property, as well as most other properties of interest, follow from the following more general statement.

Let  $(\mathbf{C}, \mathbf{C}_{\text{adm}}, \mathbf{C}^{\text{coadm}})$  and  $(\mathbf{D}, \mathbf{D}_{\text{adm}}, \mathbf{D}^{\text{coadm}})$  be two triples as above. Let  $p : \mathbf{C} \rightarrow \mathbf{D}$  be a functor which preserves admissible morphisms (resp. co-admissible morphisms) and bi-admissible cartesian squares. First of all, note:

**Exercise 1.4.2.** *The functor  $p$  induces a canonical functor*

$$(1.1) \quad p : \mathbf{Corr}(\mathbf{C}) \rightarrow \mathbf{Corr}(\mathbf{D}).$$

The fundamental fact about the construction  $\mathbf{C} \mapsto \mathbf{Corr}(\mathbf{C})$  is:

**Proposition 1.4.3** (Barwick). *If  $p$  is an inner fibration, then (1.1) is an inner fibration.*

This fact is out of our reach, but we have enough technology to check:

**Exercise 1.4.4.** *If  $p$  is a Kan fibration, then (1.1) is a Kan fibration.*

#### REFERENCES

- [1] Denis-Charles Cisinski, *Higher category theory and homotopical algebra*, Lecture notes, 2016–7, Available at <http://www.mathematik.uni-regensburg.de/cisinski/CatLR.pdf>.
- [2] Clark Barwick, *Spectral Mackey functors and equivariant algebraic K-theory (I)*.